



## Images in Clinical Rheumatology

### IgG4-related orbital pseudotumor\*

### Pseudotumor orbitario relacionado con IgG4

Rebeca Fernández Regueiro,<sup>a,\*</sup> Eva María Fonseca Aizpuru,<sup>a</sup> Carolina Estrada Menéndez,<sup>b</sup> Loreto Buznego Suárez<sup>c</sup>

<sup>a</sup> Servicio de Medicina Interna, Hospital Universitario de Cabueñas, Gijón, Asturias, Spain

<sup>b</sup> Centro de Salud de Contruces, Gijón, Asturias, Spain

<sup>c</sup> Servicio de Oftalmología, Hospital Universitario de Cabueñas, Gijón, Asturias, Spain

#### ARTICLE INFO

##### Article history:

Received 30 November 2017

Accepted 7 February 2018

Available online 5 June 2019

A 47 year-old woman with a background of total thyroidectomy from goitre with benign thyroid nodules presented with exophthalmos of several weeks onset (Fig. 1). There was no involvement of the ENT sphere, and no pulmonary, polyneuropathy, dry syndrome, arthralgias, photosensitivity or cutaneous lesions. Magnetic resonance (MR) imaging revealed a solid retrobulbar mass which surrounded the optic nerve and was of fusiform morphology measuring 24 × 9 mm (Fig. 2). We considered the need for biopsy but desisted, due to the high risk of complications. The ANAS, ANCAS, anti-DNA analysis, complement activity, proteinuria and ACE tested negative and the raising of immunoglobulin G4 860 mg/dl was remarkable (normal: 10–140). The patient was diagnosed with a possible IgG4-related inflammatory orbital disease in keeping with clinical and serological criteria. Treatment with high doses of steroids and azatioprine was initiated but suspended due to intolerance. The patient is currently being treated with mycophenolate mofetil and is responding favourably with no inflammatory activity. IgG4-related disease is a new entity which comprises several fibroinflammatory diseases which have not been previously linked to one another.<sup>1,2</sup> Clinical features are highly varied and presentation is usually subacute.<sup>3,4</sup> Ocular-orbital symptoms include: idiopathic orbital inflammation, pachymeningitis and sclerouveitis.<sup>5,6</sup> Orbital pseudotumour is an uncommon presentation of IgG4 diseases.<sup>7</sup> It is important to recognize this new disease to diagnose it, since specific treatment may prevent complications.



Fig. 1. Exophthalmos.

\* Please cite this article as: Fernández Regueiro R, Fonseca Aizpuru EM, Estrada Menéndez C, Buznego Suárez L. Pseudotumor orbitario relacionado con IgG4. Reumatol Clin. 2020;16:245–246.

\* Corresponding author.

E-mail address: [rfernandezregueiro@yahoo.es](mailto:rfernandezregueiro@yahoo.es) (R. Fernández Regueiro).



Fig. 2. MR imaging with pseudotumour.

#### Conflict of interests

The authors have no conflict of interests to declare.

#### References

1. Stone JH, Zen Y, Deshpande V. IgG4-related disease. *N Engl J Med.* 2012;366:539–51.
2. Ardila-Suarez O, Abril A, Gómez-Puerta JA. IgG4-related disease: a concise review of the current literature. *Reumatol Clin.* 2017;13:160–6.
3. López Vivancos J, Sanchez Sitges L, Herranz Perez X, Leal Bohorquez N, Simonetti S, García Cors M. IgG4-related disease with possible myocardial involvement [Article in English, Spanish]. *Reumatol Clin.* 2017, pii: S1699-258X(17)30268-1.
4. Kamisawa T, Zen Y, Pillai S, Stone JH. IgG4-related disease. *Lancet Lond Engl.* 2015;385:1460–71.
5. Mombaerts I, Bilyk JR, Rose GE, McNab AA, Fay A, Dolman PJ, et al. Consensus on diagnostic criteria of idiopathic orbital inflammation using a modified Delphi approach. *JAMA Ophthalmol.* 2017;135: 769–76.
6. Wallace ZS, Deshpande V, Stone JH. Ophthalmic manifestations of IgG4-related disease: single-center experience and literature review. *Semin Arthritis Rheum.* 2014;43:806–17.
7. Lee CS, Harocopoulos GJ, Kraus CL, Lee AY, van Stavern GP, Couch SM, et al. IgG4-associated orbital and ocular inflammation. *J Ophthalmic Inflamm Infect.* 2015;5:15.