



Original Article

The HLA-B*51 Allele is strongly associated with Behcet Disease in an Argentinean population



Sebastián Andrés Muñoz^a, Alberto Omar Orden^b, Alex Kostianovsky^c, Cecilia N. Pisoni^c, Marina Scolnik^d, Aurelia Luissi^d, Yanina Bottinelli^e, Gustavo Vijoditz^e, Mercedes Garcia^f, Claudia Pena^f, Mariana Pera^f, Oscar Rillo^g, Teresita Alvarellos^h, Luciana M. Más^h, Arturo Luis Trunzoⁱ, Alberto Allievi^{j,*}

^a Hospital General de Agudos Dr. Juan A. Fernández, Ciudad Autónoma de Buenos Aires, Argentina^b Clínica San Camilo, Ciudad Autónoma de Buenos Aires, Argentina^c Hospital Universitario CEMIC, Ciudad Autónoma de Buenos Aires, Argentina^d Hospital Italiano de Buenos Aires, Ciudad Autónoma de Buenos Aires, Argentina^e Hospital Nacional Dr. Alejandro Posadas, Ciudad del Palomar, Provincia de Buenos Aires, Argentina^f Hospital General José de San Martín, Ciudad de La Plata, Provincia de Buenos Aires, Argentina^g Hospital General de Agudos Dr. Ignacio Pirovano, Ciudad Autónoma de Buenos Aires, Argentina^h Hospital Privado Universitario de Córdoba, Ciudad de Córdoba, Provincia de Córdoba, Argentinaⁱ Instituto Nacional Central Único Coordinador de Ablación e Implante (INCUCAI), Ciudad Autónoma de Buenos Aires, Argentina^j Medicina, Universidad del Salvador, Ciudad Autónoma de Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 6 April 2018

Accepted 2 July 2018

Available online 6 August 2018

Keywords:

Systemic vasculitis

Behcet's disease

HLA-B antigen

HLA-B51 antigen

Case-control studies

Latin America

ABSTRACT

Objective: To assess the association between the HLA-B*51 allele and Behcet Disease (BD) in Argentinean patients.

Methods: We enrolled 34 consecutive Argentinean patients with definitive diagnosis of BD between October 2016 and March 2017. None of the patients had the HLA-B*51 allele determined at study entry. Unrelated controls ($n = 240$) were randomly obtained from the national cadaveric donor database. Demographic and clinical features of the patients were recorded by attending physicians through a questionnaire.

Results: Mean age of cases was 42 years old. Nineteen (55.8%) were male, and the mean age at diagnosis was 35 years old; twenty (58.8%) were Mestizos, 8 (23.5%) were Caucasian, and 6 (17.6%) were Amerindians. Thirteen (38.2%) of 34 cases were HLA-B*51 allele positive; 11 were heterozygous and 2 homozygous for the allele. Thirty-four (14.2%) of 240 controls were positive for the HLA-B*51 allele. The association between BD and HLA-B*51 allele was greater than that of control group (OR = 3.75; $p = 0.0012$).

Conclusions: The HLA-B*51 allele is strongly associated with BD in Argentinean patients. Our finding is consistent with previous studies indicating that the HLA-B*51 allele is an important susceptibility gene in BD regardless the geographical region and ethnicity.

© 2018 Elsevier España, S.L.U. and Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. All rights reserved.

El alelo HLA-B*51 se asoció fuertemente a la Enfermedad de Behcet en la población argentina

RESUMEN

Palabras clave:

Vasculitis sistémicas

Enfermedad de Behcet

Antígeno HLA-B

Antígeno HLA-B51

Estudios de casos controles

Latinoamericanos

Objetivo: Evaluar la asociación entre el alelo HLA-B*51 y la enfermedad de Behcet (EB) en pacientes argentinos.

Métodos: Incluimos en forma consecutiva 34 pacientes argentinos con diagnóstico definitivo de EB entre los meses de octubre de 2016 y marzo de 2017. Ninguno de los pacientes tenía el alelo HLA-B*51 determinado al inicio del estudio. Los controles no relacionados ($n = 240$) se obtuvieron al azar de la base nacional de datos de donantes cadávericos.

* Corresponding author.

E-mail address: beto.allievi@gmail.com (A. Allievi).

Las características demográficas y clínicas de los pacientes fueron registradas por los médicos asistentes a través de un cuestionario.

Resultados: La edad promedio de los casos fue de 42 años. Diecinueve (55,8%) fueron varones, y la edad promedio en el momento del diagnóstico fue de 35 años; 20 (58,8%) fueron mestizos, 8 (23,5%) caucásicos y 6 (17,6%) amerindios. Trece (38,2%) de los 34 casos fueron positivos para el alelo HLA-B*51; 11 de ellos fueron heterocigotas y 2 homocigotas para dicho alelo. Treinta y cuatro (14,2%) de los 240 controles fueron positivos para el alelo HLA-B*51. La asociación entre la EB y el alelo HLA-B*51 fue mayor que en el grupo control ($OR = 3,75$; $p = 0,0012$).

Conclusiones: El alelo HLA-B*51 está fuertemente asociado con la EB en pacientes argentinos. Nuestro hallazgo es consistente con estudios previos que indican que el alelo HLA-B*51 es un gen de susceptibilidad importante en la EB independientemente de la región geográfica y la etnia.

© 2018 Elsevier España, S.L.U.

y Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. Todos los derechos reservados.

Introduction

Behçet disease (BD) is a systemic inflammatory disease of unknown etiology. It is characterized by recurrent oral aphthous and genital ulcers, uveitis, and skin lesions. Vascular injury, hyper-coagulability, hyper-function of neutrophils, and autoimmune response are pathological findings of the disease.^{1,2}

BD is a complex clinical entity where genetic and environmental factors influence disease' susceptibility. Several previous studies showed that immunogenetics factors (mainly HLA class A and B) are present in the development of BD.³

HLA-B*51 allele is the most strongly associated genetic risk factor linked with BD.^{1,2,4–6} This association is present along different geographical regions and ethnic groups.⁷ So far, two studies from Mexico are the only evidence on the association between the HLA-B51 allele and BD in Latin American countries; however, the results are contradictory.^{8,9}

The aim of this study was to assess the association between BD and the HLA-B*51 allele in Argentinean patients.

Materials and Methods

Study Design and Setting

We carried out a case-control study. The researchers who participated in the study work in reference Hospitals, and they belong to the Vasculitis Study Group of the Argentinean Society of Rheumatology.

Participants

Thirty-four consecutive BD Argentinean patients were included for the study between October 2016 and March 2017. All of them had a definitive diagnosis of BD according to the International Study Group for Behçet Disease criteria, and they did not have the HLA-B*51 allele determined at study entry.¹⁰ Unrelated controls ($n = 240$), and their HLA-B*51 status, were randomly obtained from the cadaveric donor database from the “Instituto Nacional Central Único Coordinador de Ablación e Implante del Ministerio de Salud de La Nación (INCUCAI)” the official national institution for transplant allocation in Argentina.

Variables

Demographic and clinical features of the patients were recorded by attending physicians through a questionnaire. Ethnicity was defined according to definitions made by the “Grupo Latinoamericano de Estudio del Lupus (GLADEL).”¹¹

HLA Class I Typing

For cases HLA class I typing was performed using a polymerase chain reaction sequence-specific oligonucleotide (PCR-SSO) coupled with a Luminex platform using a medium resolution LABType SSO class I B locus typing test (One Lambda Inc., California, USA) according to the manufacturer's instructions. The controls were typed by a PCR with sequence-specific primers (PCR-SSP).

Statistical Analysis

Descriptive statistics including percentages, mean, standard deviation (SD), median and range were calculated. The frequencies of the alleles and genotypes among patients and controls were compared by chi-squared tests. Odds ratios and 95% confidence intervals were calculated. A p level of ≤ 0.05 was considered significant in all analysis. Statistical analysis was done using Stata 14.0 (StataCorp 2015, TX, USA).

Ethics Approval

The study was approved by all local ethical committees of the corresponding hospitals, and all the study participants gave their written informed consent to participate. The research methods were in compliance of the tenets of the Declarations of Helsinki.

Results

Demographics and Clinical Features

Main demographic characteristics are shown in Table 1. BD was slightly more prevalent in males, with a mean age at diagnosis of 35 years old. Mestizo population accounted for the majority of ethnicity, followed by White and Amerindians subjects. Most frequent clinical manifestations were oral and genital ulcers, followed by cutaneous, articular, ocular, and neurological involvement.

HLA-B*51 Gene Distribution

Out of 34 patients with BD, 13 (38.2%) of them were positive for the HLA-B*51 allele; 11 (32.3%) were heterozygous and 2 (5.8%) were homozygous. In the control group, 34 out of 240 subjects had the HLA-B*51 allele (14.2%). None of the controls were homozygous for the allele HLA-B*51. The association between BD and HLA-B*51 allele was greater than that of control group ($OR = 3.75$; 95%IC = 1.71–8.19; $p = 0.0012$) (Table 2).

Table 1

Demographic and Clinical Features of 34 Argentinean Patients With Behcet's Disease.

Features	n = 34
Female/male; n (%)	15/19 (44.12/55.88)
Age (yrs); mean (SD)	42 (13.56)
Mean age at disease onset (yrs); mean (SD)	35 (13.86)
Mean disease duration (yrs); median (range)	15.87 (0–31.75)
Ethnic distribution; n (%)	
Mestizo	20 (58.82)
White	8 (23.53)
Amerindian	6 (17.65)
Oral ulcer; n (%)	34 (100)
Cutaneous involvement; n (%)	30 (88.24)
Erythema nodosum; n (%)	12 (35.29)
Papulopustular lesions; n (%)	22 (64.71)
Articular involvement; n (%)	25 (73.53)
Genital ulcer; n (%)	23 (67.65)
Eye involvement; n (%)	22 (64.71)
Neurologic involvement; n (%)	17 (50)
Headache; n (%)	15 (44.12)
General symptoms; n (%)	10 (29.41)
Deep vein thrombosis; n (%)	5 (14.71)
Epididymitis; n (%)	2 (5.88)

SD: standard deviation; yrs: years.

Table 2

Association of HLA-B*51 in Behcet's Disease Patients and Controls.

HLA-B*51	BD patients N = 34 (%)	Controls N = 240	p	OR (95% CI)
Present	13 (38.24)	34 (14.16)		3.75
Absent	21 (61.76)	206 (85.84)	0.0012	(1.71–8.19)

BD: Behcet' disease; OR: odds ratio; CI: confidence interval.

Table 3

HLA-B*51 Allele Frequencies in Argentinean BD Patients, Mexican Mestizo BD Patients and Controls.

Population	BD patients n ^a (a ^f)	Controls n ^a (a ^f)	p	OR (95% CI)
Argentinean	15 (0.22)	34 (0.07)	0.0012	3.71 (1.89–7.26)
Mexican Mestizo ¹⁶	6 (0.09)	10 (0.05)	0.24	1.88 (0.58–5.93)

BD: Behcet' disease; OR: odds ratio; CI: confidence interval; n^a: number of alleles; a^f: allele frequencies.

Discussion

The most significant finding in our study is a strong association between HLA-B*51 allele and Behcet disease in Argentinean patients. This association is well known in patients belonging to Asian and Eurasian populations from Japan to the Middle East.⁷ However, there is less data on Caucasian patients with BD from other regions, such as North Western Europe and North America.^{12–16} Furthermore, there are two Latin American studies focusing on the association between HLA-B*5 allele and BD. Lavalle, et al. found that the HLA-B*5 allele was more frequent in Mexican Mestizo patients with BD than ethnically matched controls (70% vs. 31%, $p < 0.05$).⁸ On the other hand, Soto Vega et al., found no association between HLA-B*51 allele and BD among Mexican Mestizo patients and controls [9% vs. 5%, OR = 1.88 (95%CI = 0.58–5.93, $p = 0.24$)].⁹ Table 3 shows the HLA-B*51 allele frequencies in our BD patients and the Mexican Mestizo patients from Soto Vega study.

In our study the frequency of the HLA-B*51 allele in BD patients was 38.2% as compared to 14.2% in controls. Strikingly, our data are quite similar to the pooled prevalence estimated for HLA-B*51 allele in the Northern/Eastern regions of Europe (BD = 39%; 95%CI = 28.2–51.1 vs. controls = 11.2%; 95%CI = 8.1–15.3) and North America (BD = 34.2%; 95%CI = 6.0–80.8 vs. controls = 18%; 95%CI = 7.6–37.1), but lower than the pooled prevalence estimated

in the Southern Europe (BD = 60.6%; 95% CI = 51.9–68.7 vs. controls = 16.8%; 95% CI = 13.3–21.0) where the biggest proportion of Argentine immigration comes from.⁷ This finding could be partially explained by the fact that our studied population was predominantly Mestizo, a mixture of European (Caucasian) and native ancestors (Amerindians). Two studies showed low prevalence of HLA-B*51 in Amerindian tribes of South America (mainly Argentine natives). Fernandez-Viña et al.,¹⁷ analyzed the prevalence of HLA antigens in a Toba tribe, finding an HLA-B*51 allele frequency of 3%. Parham et al.,¹⁸ studied the prevalence of HLA antigen in a Guarani tribe, finding an HLA-B*51 allele frequency of 6.6%. Therefore, we assume that the admixture between European and South American Amerindian could explain that mixed populations have more HLA class I allele diversity than each ethnic group separately.

Limitations

Our study has limitations. Classical case-control studies designed to evaluate gene association are exposed to bias due to the existence of population substructures that cannot be certainly determined. It is possible that our cases are not representative of all BD patients from an Argentinean population as a consequence of a selection bias. Finally, as we did not have the sex and ethnicity data of the controls, we could not match both groups for these variables.

Conclusions

The HLA-B*51 allele is strongly associated with BD in Argentinean patients. Individuals carrying HLA B*51 allele have a 4-fold susceptibility risk increase of BD compared to non-carrier individuals of the allele. Our finding is consistent with previous studies indicating that the HLA-B*51 allele is an important susceptibility gene in BD regardless the geographical region and ethnicity.

Funding

This research was funded by an Argentinean Rheumatology Society grant.

Conflicts of Interest

The authors have no conflicts of interest to declare.

References

- Barnes CG, Yazici H. Behcet's syndrome. *Rheumatology*. 1999;38:1171–6.
- Sakane T, Takeno M, Suzuki N, Inaba G. Behcet disease. *N Engl J Med*. 1999;341:1284–91.
- Montes-Cano MA, Conde-Jaldón M, García-Lozano JR, Ortiz-Fernández L, Ortega-Centeno N, Castillo-Palma MJ, et al. HLA and non-HLA genes in Behcet disease: a multicentric study in the Spanish population. *Arthritis Res Ther*. 2013;15:R145.
- Ono S, Aoki K, Suguri S, Nakayama E, Itakura K. HL-A5 and Behcet's disease. *Lancet*. 1973;2:1383–4.
- Ohno S, Ohguchi M, Hirose S, Matsuda H, Wakisaka A, Aizawa M. Close association of HLA-Bw51 with Behcet's disease. *Arch Ophthalmol*. 1982;100:1455–8.
- de Menthon M, Lavalle MP, Maldini C, Guillemin L, Mahr A. HLA-B51/B5 and the risk of Behcet's disease: a systematic review and meta-analysis of case-control genetic association studies. *Arthritis Rheum*. 2009;61:1287–96.
- Verity DH, Marr JE, Ohno S, Wallace GR, Stanford MR. Behcet's disease, the Silk Road and HLA-B51: historical and geographical perspectives. *Tissue Antigens*. 1999;54:213–20.
- Lavalle C, Alarcón-Segovia D, Del Guidice-Knipping JA, Fraga A. Association of Behcet's syndrome with HLA-B5 in the Mexican mestizo population. *J Rheumatol*. 1981;8:325–7.
- Soto Vega E, García-Muñoz R, Richaud-Patin Y, et al. Class I and class II MHC polymorphisms in Mexican patients with Behcet's disease. *Immunol Lett*. 2004;93(2–3):211–5.
- Criteria for diagnosis of Behcet's disease. International Study Group for Behcet's Disease. *Lancet*. 1990;335:1078–80.
- Pons-Estel BA, Catoggio LJ, Cardiel MH, et al. The GLADEL multinational Latin American prospective inception cohort of 1.214 patients with systemic lupus

- erythematosus: ethnic and disease heterogeneity among "Hispanics". Medicine (Baltimore). 2004;83:1–17.
12. Chamberlain MA. Behçet's syndrome in 32 patients in Yorkshire. Ann Rheum Dis. 1977;36:491–9.
 13. Lehner T, Batchelor JR, Challacombe SJ, Kennedy L. An immunogenetic basis for the tissue involvement in Behçet's syndrome. Immunology. 1979;37:895–900.
 14. O'Duffy JD, Taswell HF, Elveback LR. HLA-Antigens in Behçet's disease. J Rheumatol. 1976;3:1–3.
 15. Ohno S, Char DH, Kimura SJ, O'Connor GR. Studies on HLA antigens in American patients with Behçet's disease. Jpn J Ophthalmol. 1978;22:58–61.
 16. Kilmartin DJ, Finch A, Acheson RW. Primary association of HLA-B51 with Behçet's disease in Ireland. Br J Ophthalmol. 1997;81:649–53.
 17. Fernández-Viña M, Lazaro AM, Marco CY, et al. Dissimilar evolution of B-locus compared to A-locus and class II loci of the HLA region in South American Indian tribes. Tissue Antigens. 1997;50:233–50.
 18. Parham P, Arnett KL, Adams EJ, et al. Episodic evolution and turnover of HLA-B in the indigenous human populations of the Americas. Tissue Antigens. 1997;50:219–32.