




Sociedad Española
de Reumatología -
Colegio Mexicano
de Reumatología

Reumatología clínica

www.reumatologiaclinica.org



Special Article

Hierarchical nomenclature in rheumatology*

Beatriz Yoldi Muñoz,^{a,*} María Auxiliadora Martín Martínez,^b Marta Valero Expósito,^c Carlos Plana Veret,^d José Luis Andreu Sánchez,^e José Vicente Moreno Muelas^f

^a Hospital Universitario Dexeus, Quirón Salud, Barcelona, Spain

^b Unidad de Investigación, Sociedad Española de Reumatología, Spain

^c Hospital Universitario Ramón y Cajal y Hospitales HM, Madrid, Spain

^d Hospital Clínic Provincial, Barcelona, Spain

^e Hospital Universitario Puerta de Hierro, Majadahonda, Madrid, Spain

^f Hospital Vall d'Hebrón, Barcelona, Spain



ARTICLE INFO

Article history:

Received 29 November 2018

Accepted 29 November 2018

Keywords:

Rheumatology

Diagnostic techniques

Procedures therapeutics

ABSTRACT

Introduction: One of the missions of the Spanish Society of Rheumatology is to provide the necessary tools for excellence in health care. Currently, there is no reference point to quantify medical actions in this specialty, and this is imperative.

Material and method: A list of actions was drawn up and a hierarchical classification system was established by developing a complexity index, calculated based on the completion time and difficulty level of each action.

Results: The results of the Delphi method tended to the consensus opinion within a group ($\text{mean } \sigma_2 - \sigma_1 = 0.75 - 1.43 = -0.68$, $\text{mean IQR}_2 - \text{IQR}_1 = 0.8 - 1.9 = -1.1$). The values of the complexity index ranged between 48 and 465 points. Among consultation actions, those reaching the highest scores were the first inpatient visit (366) and visits to the patient's home (369). Among diagnostic techniques, biopsies were prominent, those with the highest score were: bone biopsy (465), sural nerve biopsy (416) and synovial biopsy (380). Ultrasound scan scored 204, capillaroscopy 113 and densitometry 112. Among therapeutic techniques, infiltration/arthrocentesis/articular injection in children reached the highest difficulty (388). The score for ultrasound-guided articular injection was 163. The score for clinical report on disability was 323 and expert report 370.

Conclusions: A nomenclature of 54 actions in Rheumatology was compiled. Biopsies (bone, sural nerve, synovial), inpatient visits, visits to the patient's home, infiltrations in children, and the preparation of the expert report were identified as the most complex actions. Musculoskeletal ultrasound is twice as complex as subsequent visits, capillaroscopy or bone densitometry.

© 2018 Elsevier España, S.L.U. and Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. All rights reserved.

Nomenclátor jerarquizado en reumatología

RESUMEN

Palabras clave:

Reumatología

Técnicas diagnósticas

Procedimientos terapéuticos

Introducción: Una misión de la Sociedad Española de Reumatología es aportar las herramientas necesarias para alcanzar la excelencia asistencial. En la actualidad no existe una referencia que cuantifique la complejidad de los actos médicos de esta especialidad.

Material y método: Se elaboró una relación de los actos propios del reumatólogo y se estableció un sistema de clasificación jerárquica a partir de la construcción de un índice de complejidad, calculado mediante el tiempo de realización y el grado de dificultad de cada acto.

* Please cite this article as: Yoldi Muñoz B, Martín Martínez MA, Valero Expósito M, Plana Veret C, Andreu Sánchez JL, Moreno Muelas JV. Nomenclátor jerarquizado en reumatología. Reumatol Clin. 2020;16:3-10.

* Corresponding author.

E-mail address: yoldi.ser@gmail.com (B. Yoldi Muñoz).

Resultados: Los resultados del método Delphi tendieron a una opinión grupal consensuada (media $\sigma^2 = 0,75$ - $1,43 = -0,68$, media IQR2 - IQR1 = $0,8$ - $1,9 = -1,1$). El rango de valores del índice de complejidad osciló de 48 a 465 puntos. Entre las consultas, las que alcanzaron mayor gradación fueron la primera visita al paciente hospitalizado (366) y la visita a domicilio (369). Entre las técnicas diagnósticas, destacaron las biopsias. Las que puntuaron más alto fueron: biopsia ósea (465), de nervio sural (416) y sinovial (380). La ecografía tuvo una puntuación de 204, la capilaroscopia de 113 y la densitometría de 112. Entre las técnicas terapéuticas, la máxima dificultad (388), la alcanzó la infiltración/artrocentesis/inyección articular infantil. La puntuación de la inyección articular con control ecográfico fue de 163. El informe clínico de minusvalía, 323 y el informe pericial, 370.

Conclusiones: Este trabajo ha permitido elaborar un nomenclátor de 54 actos en reumatología donde se identifican como actos más complejos la realización de biopsias (ósea, nervio sural, sinovial), la visita a paciente hospitalizado, la visita a domicilio, la infiltración infantil bajo sedación y la elaboración de un informe pericial. La ecografía osteomuscular es considerada el doble de compleja de una visita sucesiva, la capilaroscopia o la densitometría ósea.

© 2018 Elsevier España, S.L.U.

y Sociedad Española de Reumatología y Colegio Mexicano de Reumatología. Todos los derechos reservados.

Introduction

Rheumatology is the part of medicine that studies, diagnoses and treats medical diseases of the musculoskeletal system and connective tissue.¹

The rapid transformation undergone by rheumatology from the syndromes described a century ago (arthritis, arthrosis, gout) to our days has affected not only the pathologies that we can identify, but also the diagnostic techniques that have been developed and the treatments that are used. Rheumatology has clearly evolved more quickly than the resources and criteria which are allocated to our work.

One of the missions of the Sociedad Española de Reumatología (SER) is to supply all of the agents involved in treating rheumatological patients with the instruments that are necessary to improve their quality of life, ensuring they receive the best treatment. To this end it is necessary to create a list of the services that are available in excellent rheumatological care in the 21st century, while also determining their degree of complexity. This list will make it possible to offer the whole range of diagnostic-therapeutic procedures used in rheumatology, together with the standardised degree of complexity of each intervention. It will also be a fundamental instrument for minimising geographical variability in the offer of medical procedures, ensuring that excellent rheumatological care is available throughout Spanish territory.

A list of updated medical interventions agreed by rheumatology professionals and ranked according to their level of complexity would function as a reference to reduce variability in resource availability and improve the quality of care for patients with rheumatic pathology. Although several working criteria currently exist for this purpose, they are obsolete and none of them meets the needs of doctors. The nomenclature of the Organización Médica Colegial does not cover a large part of the work in the speciality of rheumatology² (Appendix B). On the other hand, the terms applied by insurance companies are very short and incomplete, and they vary widely from province to province.

There is therefore a need to update, broaden and standardise these criteria. In 2006 Olivé et al. prepared a nomenclature for the speciality of Rheumatology with the aim of validating acronyms.³ Although this work covered medical interventions, it did not take their complexity into account. This study has the aim of preparing a nomenclature of medical interventions that is based on a complexity index (CI) which includes the difficulty of the said interventions as well as how long they take to implement. The duration that is desirable in carrying out each medical intervention was stipulated in the work by Alonso et al.⁴ using the criterion of care quality. This was set in their work using a similar method to ours, that

of agreement by a panel of experts using the two-round Delphi process.

The CI was suggested as an approximate measurement of the difficulty of care, with all of the limitations that this involves, as there is no precedent at all in the literature in this field that could be used as a standardised template. Other complex elements such as doctors' experience or training, the possibility of access to technology or the cost of materials were not included, to prevent increasing variability and arbitrariness.

Methods

Group of experts

A group of experts was formed to prepare the CI, composed of a head researcher and 30 panellists. The panellists were selected on the basis of their level of knowledge of the subject and degree of collaboration, regardless of their training, their function or the level of their position. The selection criteria were years of experience and representativeness in terms of sex, geographical distribution and public and private care.

Identification and selection of rheumatological interventions

The first step towards obtaining a nomenclature of medical, diagnostic and therapeutic procedures was to identify rheumatological interventions using the studies published with this purpose^{3,4} and the opinion of the panel of experts. The study of the definition of each one of them may be consulted in Tables 1-3 of Appendix B (supplementary material available in Internet).

Data gathering

Data was gathered using questionnaires sent to the experts, and consensus was achieved using the two-round Delphi method.

Questionnaires

The questionnaires were sent to the experts by email, so that it was easy for them to answer while ensuring anonymity. Questions were prepared on the degree of difficulty of techniques and their experience in certain processes associated with rheumatological interventions. Experience was defined as their experience in performing the said technique at some time during their clinical practice. The degree of complexity was defined on a scale of from 0 to 10, where 0 was the lowest degree of complexity and 10 the highest.

Data from the first round of questions were analysed to give the general opinion. To this end the average, standard deviation (SD), the median and the interquartile range (IQR) were given for each rheumatological intervention. Once the data from the first round had been collected and analysed, a new questionnaire was designed and sent to the group of experts by email. This second questionnaire asked the experts again for their opinion on the complexity of rheumatological interventions, showing the results of the first round. The aim of this second round was to reduce the scatter of the first round opinions (measured with standard deviation and interquartile range) specifying the agreed average opinion. The experts gave their opinion, justifying it when it diverged strongly from the group view. Analysis of this second round took the same form as it had for the first. When an expert did not answer the second round of questionnaires, they were eliminated from the analysis.

Methodology for the preparation of the complexity index

To obtain the nomenclature of rheumatological procedures, a hierarchical classification system was created according to their degree of complexity and how long each one takes to perform. A complexity index was calculated as the product of the degree of complication of the procedure and the number of minutes it takes to perform. The degree of difficulty of each rheumatological intervention was obtained using the Delphi process, while the time each rheumatological intervention takes was determined by the findings published by Alonso et al.⁴

Validation of the complexity index

The complexity index was validated by a mass survey of the rheumatologists who are members of the SER, asking them about the degree to which they agreed or disagreed with the results obtained. A Likert scale from 1 to 5 was used for this, where 1 was maximum disagreement and 5 was maximum agreement. The quantitative descriptive statistics were summarised to analyse the results (the average and confidence interval, the median and interquartile range) and the frequency of the degree of agreement-disagreement with the results for each intervention.

Results

A total of 54 interventions were identified and evaluated. 93.3% (28/30) of the panellists took part in the Delphi method.

First round Delphi

Table 1 shows the results of the first Delphi round. It can be seen that the dispersion of the opinions of the 28 experts amounted to 1.4 points (SD) and 1.9 points (IQR). Only 3 interventions of the 54 had a SD of less than 1 point, and no intervention had an IQR of less than 1 point.

These results suggest the need for a new phase to reach a higher level of agreement in expert opinions of the complexity of each intervention.

Second round Delphi

After the second phase of the Delphi method, a higher level of agreement was attained in the overall opinion of the complexity of rheumatological interventions. Thus the SD of opinions was 0.75 points and the IQR was 0.8 points. Only 4 of the 54 interventions had a SD higher than 1 point, while 4 of the 54 interventions had an

IQR higher than 1 point. The results of the second round of Delphi are shown in **Table 2**.

Complexity index for medical, diagnostic and therapeutic interventions

The interventions that were evaluated scored from 48 to 465 points for complexity. When they were classified according to the type of intervention (consultation, diagnosis and therapeutic), the consultations which scored the highest were the first visit to a hospitalised patient (366) and home visits (369).

Biopsies scored the highest for diagnostic techniques: bone (465), sural nerve (416) and synovial (380) biopsy. Ultrasound scans scored 204, capillaroscopy 113 and densitometry 112.

Of the therapeutic techniques, intra-articular injection/arthrocentesis/infiltration in children under sedation all scored 388 for complexity, while ultrasound scan-controlled intra-articular injection scored 163. A clinical report of disability was agreed to score 323 on the index, and an expert report 370. The complexity index for all of the interventions evaluated is shown in **Table 3**.

Validation of the complexity index

The complexity index was sent to a total of 1144 rheumatologists for validation; 175 answered, so that the response rate amounted to 15.3%. At least 70% of the respondents agreed or agreed strongly with the complexity index obtained for each intervention. The validation results are shown in **Table 4**.

Discussion

This work made it possible to prepare an updated ranked nomenclature of the 54 medical procedures undertaken by Spanish rheumatologists in their habitual clinical practice.

The reference criterion for the whole state is the one approved by the Organización Médica Colegial,² with data from the year 2003. This only includes 20 procedures, of which 4 are hospital visits on different days after admission. This criterion contains no more than half of the interventions which are intrinsic to rheumatology. Individually, each hospital and each insurance company applies a different list of care interventions, and these lists vary widely.

Caring for patients with systemic or inflammatory pathology requires the quantification of outcome assessments to establish treatment strategies, in the search for a therapeutic objective. For decades, patient perception of their own health and how disease affects different areas of their life has been taken into account in evaluating quality of life. It is also used as a disease activity parameter. These assessments are obtained in part from questionnaires completed by patients, which are routinely applied in medical departments (Patient Reported Outcomes). Their use and application are considered to be indispensable for the standardised quantification of working in many pathologies.^{5–7} This is also the case for therapeutic decision-making and measuring the results of clinical trials, so that they have to be included as tasks which are intrinsic to the speciality. Depending on whether they consist of evaluations of quality of life, psychopathological factors, functional impact or pain, their complexity index is said to stand at from 58.5 to 69.3.

In recent decades rheumatology has seen exponential progress in new treatments. The administration of intravenous therapies such as bisphosphonates in metabolic bone diseases, biological therapies in systemic or inflammatory diseases and intravenous immunoglobulin or cyclophosphamide all form part of the usual routine in contemporary rheumatology. Due to the need to administer these drugs, rheumatology departments began to include day

Table 1

Complexity of medical, diagnostic and therapeutic interventions in rheumatology. First round Delphi.

	N	Average	SD	Median	IQR ₁	Average experience ^a
Rheumatological visits						
First visit	28	7.71	1.41	8.0	2.0	–
Second visit	28	6.96	1.29	7.5	2.0	–
Subsequent visits	28	5.57	1.14	5.0	1.3	–
Rheumatological hospital visits						
First visit to a hospitalised patient	27	8.41	1.22	9.0	1.0	–
Subsequent visits to a hospitalised patient	28	6.86	1.30	7.0	2.0	–
Discharge visit to a hospitalised patient	28	7.11	1.59	7.0	3.0	–
First referral visit to a hospitalised patient	28	7.68	1.36	8.0	2.0	–
Subsequent referral visits	28	6.32	1.33	6.0	2.3	–
Discharge referral visit	28	6.54	1.53	6.0	3.0	–
Day hospital: patient evaluation and treatment	28	6.86	1.60	7.0	2.0	–
Day hospital: patient monitoring after treatment and discharge	28	6.36	1.37	7.0	2.0	–
Day hospital: telephonic patient monitoring	28	4.96	1.23	5.0	2.0	–
Other rheumatological visits						
Patient home visits	24	8.38	1.64	9.0	3.0	–
Patient education visits	28	6.04	1.48	6.0	2.0	–
Visits for examination/medical legal report	26	8.42	1.45	8.5	1.8	–
Remote visits	28	6.61	1.95	7.0	2.3	–
Diagnostic tests						
Quality of life evaluation test	28	6.00	1.78	6.0	2.0	–
Psychopathology evaluation test	27	7.11	1.63	7.0	2.0	–
Functional impact evaluation test	28	6.14	1.80	6.0	3.0	–
Pain and fatigue evaluation test	28	5.79	1.97	6.0	3.0	–
Applying Schirmer test	28	4.79	1.89	5.0	3.0	–
Diagnostic tests						
Arthrocentesis with ultrasound control	28	7.86	1.69	8.0	2.0	8.13
Arthrocentesis with radioscopic control	26	8.50	1.63	9.0	2.0	8.00
Analysis of synovial fluid (study of crystals)	28	6.93	1.65	7.0	2.0	6.93
Synovial biopsy	26	8.92	1.29	9.0	1.0	8.36
Sural nerve biopsy	25	9.36	0.86	10.0	1.0	–
Muscle biopsy	25	9.04	1.31	9.0	1.0	7.00
Bone biopsy	25	9.32	1.22	10.0	1.0	7.00
Skin biopsy	26	7.50	1.75	8.0	1.0	6.56
Subcutaneous fat biopsy	26	7.38	1.79	7.0	2.5	7.19
Saliva gland biopsy	27	8.19	1.88	9.0	2.5	7.00
Temporal artery biopsy	25	9.16	1.14	9.0	1.0	–
Musculoskeletal ultrasound scan	27	7.52	1.34	8.0	1.0	7.65
Capillaroscopy	28	7.04	1.35	7.0	1.0	7.05
Bone densitometry	28	6.14	1.48	6.0	2.0	6.12
Carotid ultrasound scan	27	7.70	1.17	8.0	1.0	7.64
Therapeutic techniques						
Arthrocentesis and intra-articular injection	28	6.43	1.53	7.0	2.0	6.43
Ultrasound guided intra-articular injection	28	7.39	1.40	8.0	1.3	7.38
Radioscopically guided intra-articular injection	24	8.25	1.07	8.0	1.0	8.33
Soft tissue infiltration (tendon, ligament, fascia or muscle)	28	5.93	1.61	6.0	2.0	5.93
Lumbar epidural injection	26	8.12	1.37	8.0	1.0	7.50
Cervical epidural injection	24	9.25	0.94	10.0	1.3	8.00
Radioscopically guided lumbar injection	24	8.88	0.99	9.0	2.0	8.67
Radioscopically guided sacroiliac injection	24	8.83	1.01	9.0	2.0	8.80
Intra-articular injection, arthrocentesis or paediatric infiltration with sedation	26	8.65	1.32	9.0	2.0	8.62
Peripheral nerve blockage	25	8.00	1.44	8.0	2.0	7.27
Radiosynovectomy without radioscopic control (knee)	27	7.52	1.55	8.0	1.5	7.31
Radiosynovectomy with radioscopic or ultrasound control	26	8.19	1.17	8.0	2.0	8.25
Intra-articular biological implant injection	26	7.65	1.85	8.0	2.8	6.77
Treatment of soft tissue injuries. Immobilisation	26	6.42	1.65	6.5	1.8	6.41
Ultrasound guided calcification washing	16	8.13	1.15	8.0	2.0	7.83
PRP treatment with ultrasound-guided extraction and centrifuging	17	8.12	1.32	8.0	1.0	8.33
Reports						
Clinical disability report	28	7.82	1.25	8.0	2.0	–
Expert report	26	8.69	1.35	9.0	2.0	–
Overall	28	1.43		1.9		

SD: standard deviation; PRP: platelet-rich plasma; IQR: interquartile range [percentile₂₅–percentile₇₅].^a Complexity evaluated by the doctors who reported experience with this intervention.

hospital care departments, and this is now habitual. They also include the administration of diagnostic or therapeutic techniques, as well as support for clinical trials. 77.7% of rheumatologists have a day hospital (unpublished SER data). In the VALORA study, 91%

of Day Hospital Units use integrated visits that cover clinical evaluation, analysis and procedures. The average number of treatments carried out in these units by rheumatology amounted to 748.3, while it totalled 1126.73 in complex hospitals.⁸ It was found that

Table 2

Complexity of medical, diagnostic and therapeutic interventions in rheumatology. Second round Delphi.

	N	Average	SD	Median	IQR ₁	Average experience ^a
Rheumatological visits						
First visit	27	7.74	.71	8.0	1.0	–
Second visit	27	6.74	.81	7.0	1.0	–
Subsequent visits	27	5.78	.70	6.0	1.0	–
Rheumatological hospital visits						
First visit to a hospitalised patient	27	8.56	.58	9.0	1.0	–
Subsequent visits to a hospitalised patient	27	7.04	.65	7.0	.0	–
Discharge visit to a hospitalised patient	27	7.30	.91	8.0	1.0	–
First referral visit to a hospitalised patient	27	7.78	.80	8.0	1.0	–
Subsequent referral visits	27	6.41	.84	6.0	1.0	–
Discharge referral visit	27	6.48	.89	6.0	1.0	–
Day hospital: patient evaluation and treatment	26	6.92	.84	7.0	.0	–
Day hospital: patient monitoring after treatment and discharge	27	6.33	1.07	7.0	1.0	–
Day hospital: telephonic patient monitoring	26	5.04	1.11	5.0	1.0	–
Other rheumatological visits						
Patient home visits	26	8.62	.64	9.0	1.0	–
Patient education visits	25	6.00	.87	6.0	.0	–
Visits for examination/medical legal report	26	8.58	.90	9.0	1.0	–
Remote visits	26	6.77	.82	7.0	1.0	–
Diagnostic tests						
Quality of life evaluation test	27	5.85	.66	6.0	1.0	–
Psychopathology evaluation test	27	6.93	.83	7.0	1.5	–
Functional impact evaluation test	27	6.00	.78	6.0	.5	–
Pain and fatigue evaluation test	27	5.78	.93	6.0	1.0	–
Applying Schirmer test	25	4.84	.99	5.0	2.0	–
Diagnostic tests						
Arthrocentesis with ultrasound control	27	8.30	.47	8.0	1.0	8.27
Arthrocentesis with radioscopic control	25	8.40	.50	8.0	1.0	8.33
Analysis of synovial fluid (study of crystals)	27	7.00	.68	7.0	.0	6.96
Synovial biopsy	26	9.08	.56	9.0	.0	8.89
Sural nerve biopsy	25	9.48	.51	9.0	1.0	–
Muscle biopsy	25	8.96	.89	9.0	.0	8.00
Bone biopsy	25	9.36	.64	9.0	1.0	10.00
Skin biopsy	27	7.44	1.01	8.0	1.0	6.88
Subcutaneous fat biopsy	25	7.48	.92	8.0	1.0	7.46
Saliva gland biopsy	27	7.93	1.14	8.0	2.0	6.83
Temporal artery biopsy	25	9.36	.49	9.0	1.0	–
Musculoskeletal ultrasound scan	26	7.85	.46	8.0	.0	7.91
Capillaroscopy	27	7.19	.48	7.0	.0	7.20
Bone densitometry	27	6.04	.71	6.0	.0	6.07
Carotid ultrasound scan	25	7.92	.57	8.0	.0	8.00
Therapeutic techniques						
Arthrocentesis and intra-articular injection	27	6.48	.85	7.0	1.0	6.50
Ultrasound guided intra-articular injection	27	7.74	.66	8.0	1.0	7.86
Radioscopically guided intra-articular injection	24	8.21	.72	8.0	1.0	8.20
Soft tissue infiltration (tendon, ligament, fascia or muscle)	27	5.85	.91	6.0	.0	5.84
Lumbar epidural injection	25	8.48	.77	8.0	1.0	8.18
Cervical epidural injection	25	9.24	.52	9.0	1.0	8.00
Radioscopically guided lumbar injection	25	9.00	.76	9.0	.0	9.00
Radioscopically guided sacroiliac injection	25	9.00	.65	9.0	.0	9.00
Intra-articular injection, arthrocentesis or paediatric infiltration with sedation	25	9.16	.62	9.0	1.0	9.10
Peripheral nerve blockage	25	8.36	.70	8.0	1.0	8.13
Radiosynovectomy without radioscopic control (knee)	26	8.04	.87	8.0	2.0	7.83
Radiosynovectomy with radioscopic or ultrasound control	25	8.60	.76	9.0	1.0	9.00
Injection of intra-articular biological implants	24	7.25	.90	7.0	1.0	7.30
Treatment of soft tissue injuries. Immobilisation	26	6.46	.86	6.0	1.0	6.60
Ultrasound guided calcification washing	24	8.46	.51	8.0	1.0	8.40
PRP treatment with ultrasound-guided extraction and centrifuging	23	8.00	.67	8.0	.0	–
Reports						
Clinical disability report	27	7.81	.68	8.0	.5	–
Expert report	27	8.96	.59	9.0	.0	–
Overall	27		.75		.8	

SD: standard deviation; PRP: platelet-rich plasma; IQR: interquartile range [percentile₂₅–percentile₇₅].^a Complexity evaluated by the doctors who reported experience with this intervention.

work in these units should be updated and included within our nomenclature, and it was evaluated with a resulting complexity index of from 159 to 114, depending on whether it was the first or a post-treatment visit.

Rheumatology has progressed thanks to the inclusion of new treatments, and another major revolution has occurred in the use of new diagnostic and therapeutic tools and techniques such as capillaroscopy, which are intrinsic to the speciality and in widespread

Table 3

Complexity of medical, diagnostic and therapeutic interventions in rheumatology.

	Time (min.)	Complexity	Index (Complexity*time)
<i>Rheumatological visits</i>			
First visit	29.7	7.74	229.9
Second visit	18.7	6.74	126.0
Subsequent visits	17.4	5.78	100.6
<i>Rheumatological hospital visits</i>			
First visit to a hospitalised patient	42.8	8.56	366.4
Subsequent visits to a hospitalised patient	17.5	7.04	123.2
Discharge visit to a hospitalised patient	41.3	7.30	301.5
First referral visit to a hospitalised patient	37.3	7.78	290.2
Subsequent referral visits	22.0	6.41	141.0
Discharge referral visit	23.2	6.48	150.3
Day hospital: patient evaluation and treatment	23.0	6.92	159.2
Day hospital: patient monitoring after treatment and discharge	18.1	6.33	114.6
Day hospital: telephonic patient monitoring	16.1	5.04	81.1
<i>Other rheumatological visits</i>			
Patient home visits	42.8	8.62	368.9
Patient education visits	17.4	6.00	104.4
Visits for examination/medical legal report	29.7	8.58	254.8
Remote visits	16.1	6.77	109.0
<i>Diagnostic tests</i>			
Quality of life evaluation test	10.0	5.85	58.5
Psychopathology evaluation test	10.0	6.93	69.3
Functional impact evaluation test	10.0	6.00	60.0
Pain and fatigue evaluation test	10.0	5.78	57.8
Applying Schirmer test	10.0	4.84	48.4
<i>Diagnostic tests</i>			
Arthrocentesis with ultrasound control	24.3	8.30	201.7
Arthrocentesis with radioscopic control	32.0	8.40	268.8
Analysis of synovial fluid (study of crystals)	16.7	7.00	116.9
Synovial biopsy	41.8	9.08	379.5
Sural nerve biopsy	43.9	9.48	416.2
Muscle biopsy	42.1	8.96	377.2
Bone biopsy	49.7	9.36	465.2
Skin biopsy	24.1	7.44	179.3
Subcutaneous fat biopsy	19.1	7.48	142.9
Saliva gland biopsy	26.7	7.93	211.7
Temporal artery biopsy	38.6	9.36	361.3
Musculoskeletal ultrasound scan	26.0	7.85	204.1
Capillaroscopy	15.7	7.19	112.9
Bone densitometry	18.5	6.04	111.7
Carotid ultrasound scan	26.0	7.92	205.9
<i>Therapeutic techniques</i>			
Arthrocentesis and intra-articular injection	12.4	6.48	80.4
Ultrasound guided intra-articular injection	21.1	7.74	163.3
Radioscopically guided intra-articular injection	25.2	8.21	206.9
Soft tissue infiltration (tendon, ligament, fascia or muscle)	12.3	5.85	72.0
Lumbar epidural injection	27.6	8.48	234.0
Cervical epidural injection	33.4	9.24	308.6
Radioscopically guided lumbar injection	34.4	9.00	309.6
Radioscopically guided sacroiliac injection	33.1	9.00	297.9
Intra-articular injection, arthrocentesis or paediatric infiltration with the need for sedation	42.3	9.16	387.5
Peripheral nerve blockage	31.6	8.36	264.2
Radiosynovectomy without radioscopic control (knee)	25.0	8.04	201.0
Radiosynovectomy with radioscopic or ultrasound control	35.2	8.60	302.7
Injection of intra-articular biological implants	12.4	7.25	89.9
Treatment of soft tissue injuries. Immobilisation	25.0	6.46	161.5
Ultrasound guided calcification washing	35.0	8.46	296.1
PRP treatment with ultrasound-guided extraction and centrifuging	30.0	8.00	240.0
<i>Reports</i>			
Clinical disability report	41.3	7.81	322.6
Expert report	41.3	8.96	370.0

PRP: platelet-rich plasma.

* Likert scale distribution grouped as % of disagreement (1 and 2), % of indifferent (3) and % of agreement (4 and 5).

use. In fact, 72.1% of rheumatologists have direct access to this technique (unpublished SER data) together with, and more recently, ultrasound scan imaging.

Ultrasound imaging really has become a highly useful complementary form of examination for rheumatologists. It permits the harmless, static and dynamic evaluation of joint, muscle, tendon

and ligament pathologies, as well as compression and vascular neuropathies. Given its usefulness, the SER founded an ultrasound imaging school in 1996. This has a teaching staff of 28 experts, and they have taught 420 courses to date, training more than 1000 rheumatologists. 71.7% of rheumatologists had their own ultrasound scan device in 2014 (unpublished SER data). Ultrasound

Table 4
Validation of the complexity index.

	Average (CI 95%)	Median (IQR)	Response distribution (%) ^a
<i>Rheumatological visits</i>			
First visit	4.15 (4.1, 4.2)	4.0 (4.0, 5.0)	3.4/10.3/86.3
Second visit	3.91 (3.9, 4.0)	4.0 (4.0, 4.0)	4.6/20.0/75.4
Subsequent visits	3.90 (3.9, 3.9)	4.0 (3.0, 5.0)	9.1/18.9/72.0
<i>Rheumatological hospital visits</i>			
First visit to a hospitalised patient	4.44 (4.4, 4.5)	5.0 (4.0, 5.0)	1.7/6.9/91.4
Subsequent visits to a hospitalised patient	4.09 (4.0, 4.1)	4.0 (4.0, 5.0)	3.4/13.1/83.4
Discharge visit to a hospitalised patient	4.04 (4.0, 4.1)	4.0 (4.0, 5.0)	6.3/13.1/80.6
First referral visit to a hospitalised patient	4.04 (4.0, 4.1)	4.0 (4.0, 5.0)	4.6/13.9/81.5
Subsequent referral visits	3.95 (3.9, 4.0)	4.0 (3.0, 5.0)	6.9/19.5/73.6
Discharge referral visit	3.90 (3.9, 3.9)	4.0 (3.0, 4.0)	7.5/17.9/74.6
Day hospital: patient evaluation and treatment	3.90 (3.9, 3.9)	4.0 (3.0, 5.0)	6.9/22.0/71.1
Day hospital: patient monitoring after treatment and discharge	3.93 (3.9, 4.0)	4.0 (3.0, 5.0)	8.1/19.8/72.1
Day hospital: telephonic patient monitoring	4.02 (4.0, 4.1)	4.0 (3.0, 5.0)	4.7/21.9/73.4
<i>Other rheumatological visits</i>			
Patient home visits	4.12 (4.1, 4.2)	4.0 (3.0, 5.0)	5.5/22.0/72.6
Patient education visits	3.87 (3.8, 3.9)	4.0 (3.0, 5.0)	9.5/22.5/68.0
Visits for examination/medical legal report	4.30 (4.3, 4.3)	4.0 (4.0, 5.0)	4.1/8.9/87.0
Remote visits	3.83 (3.8, 3.9)	4.0 (3.0, 5.0)	10.7/17.8/71.6
<i>Diagnostic tests</i>			
Quality of life evaluation test	4.01 (4.0, 4.1)	4.0 (3.8, 5.0)	8.1/16.9/75.0
Psychopathology evaluation test	3.76 (3.7, 3.8)	4.0 (3.0, 4.0)	7.6/28.8/63.5
Functional impact evaluation test	3.94 (3.9, 4.0)	4.0 (4.0, 5.0)	5.8/18.6/75.6
Pain and fatigue evaluation test	3.87 (3.8, 3.9)	4.0 (3.0, 4.5)	7.0/22.2/70.8
Applying Schirmer test	3.92 (3.9, 4.0)	4.0 (3.0, 5.0)	10.0/19.4/70.6
<i>Diagnostic tests</i>			
Arthrocentesis with ultrasound control	4.31 (4.3, 4.4)	5.0 (4.0, 5.0)	5.2/8.7/86.1
Arthrocentesis with radioscopic control	4.24 (4.2, 4.3)	4.0 (4.0, 5.0)	5.3/10.7/84.0
Analysis of synovial fluid (study of crystals)	4.01 (4.0, 4.1)	4.0 (4.0, 5.0)	8.6/13.8/77.6
Synovial biopsy	4.65 (4.6, 4.7)	5.0 (4.0, 5.0)	1.8/4.8/93.5
Sural nerve biopsy	4.67 (4.6, 4.7)	5.0 (5.0, 5.0)	1.8/5.4/92.9
Muscle biopsy	4.59 (4.6, 4.6)	5.0 (4.0, 5.0)	2.4/7.2/90.4
Bone biopsy	4.69 (4.7, 4.7)	5.0 (5.0, 5.0)	2.4/3.6/94.0
Skin biopsy	4.23 (4.2, 4.3)	4.0 (4.0, 5.0)	5.4/10.7/83.9
Subcutaneous fat biopsy	4.16 (4.1, 4.2)	4.0 (4.0, 5.0)	4.7/15.9/79.4
Saliva gland biopsy	4.08 (4.0, 4.1)	4.0 (4.0, 5.0)	7.0/13.4/79.7
Temporal artery biopsy	4.64 (4.6, 4.7)	5.0 (4.0, 5.0)	1.8/2.4/95.8
Musculoskeletal ultrasound scan	4.16 (4.1, 4.2)	4.0 (4.0, 5.0)	4.6/13.9/81.5
Capillaroscopy	4.10 (4.1, 4.1)	4.0 (4.0, 5.0)	7.5/12.6/79.9
Bone densitometry	4.11 (4.1, 4.2)	4.0 (4.0, 5.0)	9.5/14.2/76.3
Carotid ultrasound scan	4.20 (4.2, 4.2)	4.0 (4.0, 5.0)	4.8/9.6/85.5
<i>Therapeutic techniques</i>			
Arthrocentesis and intra-articular injection	3.91 (3.9, 4.0)	4.0 (3.0, 5.0)	13.8/13.2/73.0
Ultrasound guided intra-articular injection	4.13 (4.1, 4.2)	4.0 (4.0, 5.0)	6.9/10.4/82.7
Radioscopically guided intra-articular injection	4.18 (4.1, 4.2)	4.0 (4.0, 5.0)	6.5/8.2/85.3
Soft tissue infiltration (tendon, ligament, fascia or muscle)	3.93 (3.9, 4.0)	4.0 (3.0, 5.0)	11.5/16.1/72.4
Lumbar epidural injection	4.33 (4.3, 4.4)	5.0 (4.0, 5.0)	7.1/6.5/86.3
Cervical epidural injection	4.63 (4.6, 4.7)	5.0 (4.0, 5.0)	1.8/4.8/93.4
Radioscopically guided lumbar injection	4.52 (4.5, 4.6)	5.0 (4.0, 5.0)	3.0/5.4/91.6
Radioscopically guided sacroiliac injection	4.57 (4.5, 4.6)	5.0 (4.0, 5.0)	1.8/5.4/92.8
Intra-articular injection, arthrocentesis or paediatric infiltration with sedation	4.62 (4.6, 4.7)	5.0 (4.0, 5.0)	2.4/3.6/94.0
Peripheral nerve blockage	4.43 (4.4, 4.5)	5.0 (4.0, 5.0)	1.8/7.8/90.4
Radiosynovectomy without radioscopic control (knee)	4.27 (4.2, 4.3)	4.0 (4.0, 5.0)	4.2/11.4/84.3
Radiosynovectomy with radioscopic or ultrasound control	4.33 (4.3, 4.4)	4.5 (4.0, 5.0)	3.6/9.0/87.3
Intra-articular biological implant injection	4.14 (4.1, 4.2)	4.0 (4.0, 5.0)	5.4/14.4/80.2
Treatment of soft tissue injuries. Immobilisation	4.13 (4.1, 4.2)	4.0 (4.0, 5.0)	5.4/13.9/80.7
Ultrasound guided calcification washing	4.35 (4.3, 4.4)	4.5 (4.0, 5.0)	3.0/7.7/89.3
PRP treatment with ultrasound-guided extraction and centrifuging	4.22 (4.2, 4.3)	4.0 (4.0, 5.0)	4.2/13.8/82.0
<i>Reports</i>			
Clinical disability report	4.04 (4.0, 4.1)	4.0 (4.0, 5.0)	8.2/11.7/80.1
Expert report	4.47 (4.4, 4.5)	5.0 (4.0, 5.0)	3.6/5.3/91.1

CI: confidence interval; PRP: platelet-rich plasma; IQR: interquartile range.

^a Likert scale distribution grouped as % of disagreement (1 and 2), % of indifferent (3) and % of agreement (4 and 5).

imaging has attained a position of honour in the current nomenclature of the speciality. It is an effective diagnostic tool as it offers a good approximate view of lesion anatomy and inflammatory activity by using a scale of greys⁹ and the Doppler signal,^{10,11} respectively. It is a technique that supports other imaging tests for ultrasound-guided intra-articular contrast injection. For

treatment, the spread of ultrasound imaging has led rheumatologists to adopt new interventionist techniques that can be undertaken with greater precision. These include blockages, arthrocentesis or infiltrations and ultrasound-guided treatments. These processes were graded with a complexity index of up to 387, in the case of paediatric joint puncture with sedation.

Although some techniques such as synovial, nerve, saliva gland, temporal artery or bone biopsy are not exclusive to rheumatologists, it should not be forgotten that they are habitually performed in some rheumatology units. Nor should it be forgotten that in other rheumatology units, rheumatologists themselves control pain by using more specific interventionist techniques such as epidural infiltrations.¹² The interventionist techniques defined as the most complex were biopsies, particularly bone biopsy.

Densitometry has been validated as a bone density quantification technique¹³ and its result may form part of the international indexes used to calculate the risk of fracture.¹⁴ Rheumatologists perform this test or evaluate its results regularly in their work for the diagnosis and treatment of osteoporosis. In fact, 20% of rheumatology departments have their own densitometer (unpublished SER data) so that we agree that they should be present in all lists of medical procedures concerning our speciality, with a complexity rating somewhat higher than that of a successive visit (111 vs 100).

Rheumatic diseases are one of the main causes of disability and disease burden worldwide. Data on the impact of musculoskeletal diseases on health indicate that almost 1 in 4 Spanish people have a rheumatic disease; neck pain and lumbago were the chief causes of disability in Spain in 2016.^{15,16} This justifies the frequency with which rheumatologists are asked for reports in their clinical practice that officially justify some type of disability or handicap. Moreover, rheumatologists may intervene as experts or prepare special expert reports as a part of their work outside care. Such reports were evaluated as having a complexity index of 322 and 370.

This work supplies the first ranked and agreed nomenclature of a medical speciality undertaken by a state scientific society. A tool has been constructed to improve patient care and minimise geographical variations in all areas of care.

Financing

This work was financed by the Sociedad Española de Reumatología from the assignment for Services to Members and Private Practice.

Conflict of interests

The authors have no conflict of interests to declare.

Acknowledgements

We would like to thank the panel of experts (Dr. Susana Gerechter, Dr. Juan Carlos Duró, Dr. Antonio Gómez Centeno, Dr. María Rocío González, Dr. Javier Calvo Catalá, Dr. José Ivorra, Dr. Rosa Roselló, Dr. Manuel Tenorio, Dr. Paloma Vela, Dr. Encarna Pagán, Dr. Montserrat Romera, Dr. Manuel Acasuso, Dr. Georgina Salvador, Dr. Ana Urruticoechea, Dr. Marta Larrosa, Dr. Mónica Vázquez, Dr. Cayetano Alegre, Dr. José Luis Guerra, Dr. Ana Cruz Valenciano, Dr. Emilio Martín Mola, Dr. Jesús Tornero, Dr. Javier del

Pino, Dr. Julio Medina, Dr. José Manuel Rodríguez Heredia, Dr. Eugenio Chamizo, Dr. Miguel Ángel Abad, Dr. María Luz García Vivar, Dr. Santiago Muñoz Fernández, Dr. Sara Manrique Arija and Dr. María Alcalde) for their collaboration in this project, together with all of the SER staff for their professionalism and dedication.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.reumae.2018.11.018.

References

1. Qué es la reumatología-Inforeuma. Quiénes somos. Octubre 2018. Available from: <https://inforeuma.com> [accessed 1.10.18].
2. CGCOM. Nomenclátor. Available from: <https://www.cgcom.es/sites/default/files/nomenclator.pdf> [accessed 9.10.18].
3. Olivé A, Fernández-Sueiro JL, Díaz-Jouanen E. Nomenclátor, lista de abreviaturas o acrónimos y valores de referencia de laboratorio. Reumatol Clin. 2005;1:59-75, [http://dx.doi.org/10.1016/S1699-258X\(05\)72716-9](http://dx.doi.org/10.1016/S1699-258X(05)72716-9). Núm.1.
4. Alonso A, Vidal J, Tornero J, Carbonell J, Lázaro P, Aguilar D. Estándares de calidad asistencial en reumatología. Reumatol Clin. 2007;3:218-25, [http://dx.doi.org/10.1016/S1699-258X\(07\)73690-2](http://dx.doi.org/10.1016/S1699-258X(07)73690-2). Núm.5.
5. Torre-Alonso JC, Queiro R, Comellas M, Lizán L, Blanch C. Patient-reported outcomes in European spondyloarthritis patients: a systematic review of the literature. Patient Prefer Adherence. 2018;12:733-47, <http://dx.doi.org/10.2147/PPA.S162420>, eCollection 2018.
6. Hiligsmann M, Rademacher S, Kaal KJ, Bansback N, Harrison M. The use of routinely collected patient-reported outcome measures in rheumatoid arthritis. Semin Arthritis Rheum. 2018, <http://dx.doi.org/10.1016/j.semarthrit.2018.03.006>, pii:S0049-0172(17)30577-2.
7. Robson JC, Dawson J, Doll H, Cronholm PF, Milman N, Kellom K, et al. Validation of the ANCA-associated vasculitis patient-reported outcomes (AAV-PRO) questionnaire. Ann Rheum Dis. 2018, <http://dx.doi.org/10.1136/annrheumdis-2017-212713>, pii:annrheumdis-2017-212713, PMID: 29695498.
8. Hernández Miguel MV, Martín Martínez MA, Corominas H, Sánchez-Piedra C, Sanmartí R, Fernández Martínez C, et al. Variability in Rheumatology day care hospitals in Spain: VALORA study. Reumatol Clin. 2017;13:10-6.
9. Kane D, Balint PV, Sturrock R, Grassi W. Musculoskeletal ultrasound – a state of the art review in rheumatology. Part 1: Current controversies and issues in the development of musculoskeletal ultrasound in rheumatology. Rheumatology. 2004;43:823-8. Epub 2004 May 11.
10. Joshua F, Edmonds J, Lassere M. Power Doppler ultrasound in musculoskeletal disease: a systematic review. Semin Arthritis Rheum. 2006;36:99-108.
11. Porta F, Radunovic G, Vlad V, Micu MC, Nestorova R, Petranova T, et al. The role of Doppler ultrasound in rheumatic diseases. Rheumatology (Oxford). 2012;51:976-82, <http://dx.doi.org/10.1093/rheumatology/ker433>. Publicación electrónica 16 Ene 2012.
12. Ramírez Huaranga MA, Plasencia Ezaine ÁE. Reumatología intervencionista: una asignatura pendiente. Reumatol Clin. 2018;14:2-3.
13. Roux C, Briot K. Current role for bone absorptiometry. Joint Bone Spine. 2017;84:35-7, <http://dx.doi.org/10.1016/j.jbspin.2016.02.032>. Publicación electrónica 6 Jun 2016.
14. Lewiecki EM, Compston JE, Miller PD, Adachi JD, Adams JE, Leslie WD, et al. Official Positions for FRAX® Bone Mineral Density and FRAX® simplification from Joint Official Positions Development Conference of the International Society for Clinical Densitometry and International Osteoporosis Foundation on FRAX®. J Clin Densitom. 2011;14:226-36.
15. Carmona L, Ballina J, Gabriel R, Laffon A, et al. The burden of musculoskeletal diseases in the general population of Spain: results from a national survey. Ann Rheum Dis. 2001;60:1040-5.
16. Soriano JB, Rojas-Rueda D, Alonso J, Antó J, Cardona P, Fernández E, et al. La carga de enfermedad en España: resultados del Estudio de la Carga Global de las Enfermedades 2016. Med Clin (Barc). 2017, <http://dx.doi.org/10.1016/j.medcli.2018.05.011>.