

## Images in Clinical Rheumatology

### Poor Progression After Proximal Humerus Fracture<sup>☆</sup>

#### Evolución tórpida tras fractura humeral proximal

Antonio Jiménez-Martín,<sup>a,\*</sup> Francisco Javier Santos-Yubero,<sup>a</sup> Laura Madrigal-Cortés,<sup>b</sup> Santiago Pérez-Hidalgo<sup>a</sup>

<sup>a</sup> Servicio de Cirugía Ortopédica y Traumatología, Hospital FREMAP Sevilla, Sevilla, Spain

<sup>b</sup> Servicio de Medicina Interna, Hospital FREMAP Sevilla, Sevilla, Spain

#### Introduction

Proximal humeral osteosynthesis is not without complications (45%<sup>1</sup>). Infection has a frequency of 0%–4%.<sup>2,3</sup> Our objective was to report a case of osteomyelitis, after osteosynthesis and review current treatments.

#### Case Presentation

The patient was a 54-year-old male with liver disease, allergic to penicillin, presenting a proximal humeral fracture with a cervicodiaphysal angle of 80° and a trochiter ascent of 11 mm. Pl osteosynthesis was performed with a Philos plate (Synthes®) (Fig. 1). The patient began rehabilitation, with a limited arc (60° of active abduction and 50° of anterior active flexion), fever and wound seroma (ultrasound size: 25 mm × 12 mm), at 2 months and 22 days after the intervention, requiring 2 surgical debridements and the taking of cultures and antibiograms.

#### Diagnosis

Proximal humeral osteomyelitis with stage IV B Cierny–Mader,<sup>4</sup> due to coagulase-negative *Staphylococcus*. Culture antibiogram indicated susceptibility to penicillin, cephotaxime, erythromycin, tetracycline, levofloxacin, and vancomycin cotrimoxazole. Scintigraphic uptake (Fig. 2).

#### Progression

Given the pathogenic agent, the clinical progression and the increased erythrocyte sedimentation rate (ESR: 140 mm/h), we removed the material and put a spacer cemented with gentamicin



**Fig. 1.** Osteosynthesis with plate and screws. Note the impossibility of correction of the initial cervicodiaphysal angle, worsening the mobile arc and predisposing pseudoarthrosis.

(Fig. 3). We postsurgically started (intravenous (iv)) clindamycin. The patient was under treatment with antibiotics PO: levofloxacin 500 mg 1/24 h vo rifampin 600 mg and 1/24 h PO for the first 3 months following surgery (considering the sensitivity of the pathogen and the patient's allergy to penicillin). Six months after

<sup>☆</sup> Please cite this article as: Jiménez-Martín A, et al. Evolución tórpida tras fractura humeral proximal. Reumatol Clin. 2012. <http://dx.doi.org/10.1016/j.reuma.2012.04.005>.

\* Corresponding author.

E-mail address: antonio.jimenez10@hotmail.com (A. Jiménez-Martín).




Fig. 2. Scintigraphic uptake in the right proximal humerus.



Fig. 3. Image of the spacer positioned and centered in height relative to the glenoid.

intervention the infection was controlled (ESR: 17 mm/h), with an arc of 100° (flexion/active abduction). The patient refused to undergo withdrawal of the spacer. One year after surgery, the infection remained controlled, with a similar arc.

## Discussion

Athwal et al.<sup>1</sup> recommend 3.3 surgical lavages to eradicate the infection, achieving 53 points in the *American Shoulder and Elbow Surgeons Score* (ASES), but recurrences of 50%.<sup>5</sup> Seitz et al.<sup>6</sup> recommend removing the material, placing a spacer with antibiotics and arthroplasty (although 33.3%<sup>3</sup> of patients refuse arthroplasty). Spacers improve the VAS (8.4–0.5 points), for forward flexion (65–110°) and the ASES Score (16–74 points).<sup>3</sup> Themistocleous et al.<sup>7</sup> recommend keeping the spacer, eradicating osteomyelitis in 81.8% of test with a *Disabilities of Arm, Shoulder and Hand* (DASH) score of 37.5 points. Stine et al.<sup>8</sup> maintain spacers if comorbidity is present, although controlling renal function, due to the risk of kidney failure.<sup>3,9</sup>

In conclusion, in humeral osteomyelitis cases, placement of a cemented spacer may be an effective treatment for infection control.

## Ethical disclosures

**Protection of human and animal subjects.** The authors declare that the procedures followed were in accordance with the regulations of the responsible Clinical Research Ethics Committee and in accordance with those of the World Medical Association and the Helsinki Declaration.

**Confidentiality of Data.** The authors declare that no patient data appears in this article.

**Right to privacy and informed consent.** The authors have obtained the informed consent of the patients and /or subjects mentioned in the article. The author for correspondence is in possession of this document.

## References

1. Athwal GS, Sperling JW, Rispoli DM, Cofield RH. Acute deep infection after surgical fixation of proximal humeral fractures. *J Shoulder Elbow Surg.* 2007;16:408–12.
2. Smith AM, Mardones RM, Sperling JW, Cofield RH. complications of operatively treated proximal humeral fractures. *Early J Shoulder Elbow Surg.* 2007;16:14–24.
3. Coffey MJ, Ely EE, Crosby LA. Treatment of glenohumeral sepsis with a commercially produced antibiotic-impregnated cement spacer. *J Shoulder Elbow Surg.* 2010;19:868–73.
4. Cierny 3rd G, Mader JT, Penninck JJ. A clinical staging system for adult osteomyelitis. *Clin Orthop Relat Res.* 2003;414:7–24.
5. Loebenberg MI, Zuckerman JD. An articulating interval spacer in the treatment of an infected total shoulder arthroplasty. *J Shoulder Elbow Surg.* 2004;13:476–8.
6. Seitz Jr WH, Damacén H. Staged exchange arthroplasty for shoulder sepsis. *J Arthroplasty.* 2002;17 Suppl. 1:36–40.
7. Themistocleous G, Zalavras C, Stine I, Zachos V, Itamura J. Prolonged implantation of an antibiotic cement spacer for management of shoulder sepsis in compromised patients. *J Shoulder Elbow Surg.* 2007;16:701–5.
8. Stine IA, Lee B, Zalavras CG, Hatch 3rd G, Itamura JM. Management of chronic shoulder infections utilizing a fixed articulating antibiotic-loaded spacer. *J Shoulder Elbow Surg.* 2010;19:739–48.
9. Dovas S, Liakopoulos V, Papatheodorou L, Chronopoulou I, Papavasiliou V, Atmatzidis E. Acute renal failure after antibiotic-impregnated bone cement treatment of an infected total knee arthroplasty. *Clin Nephrol.* 2008;69:207–12.