



Case Report

Peripheral Spondyloarthritis in a Patient With Noonan's Syndrome[☆]



Lina Maria Saldarriaga Rivera,* Elisa Fernandes de Melo, Priscilla Damião Araujo, Nelson Araujo Silva Filho, Luis Alberto Delgado Quiroz, Blanca Elena Rios Gomes Bica

Servicio de Reumatología. Hospital Universitario Clementino Fraga Filho, Universidad Federal de Río de Janeiro (HUCFF-UFRJ), Rio de Janeiro, Brazil

ARTICLE INFO

Article history:

Received 4 July 2013

Accepted 2 October 2013

Available online 12 March 2014

Keywords:

Noonan syndrome
Spondyloarthritis
Genetic syndrome
Arthritis

ABSTRACT

Noonan's syndrome is an autosomal dominant genetic disorder with high phenotypic variability, characterized mainly by facial dysmorphism, congenital heart disease and short stature. We describe the case of a male patient diagnosed with Noonan's syndrome and peripheral spondyloarthritis, a previously undescribed association in the literature.

© 2013 Elsevier España, S.L.U. All rights reserved.

Palabras clave:

Síndrome de Noonan
Espondiloartritis
Síndrome genético
Artritis

Presentación de espondiloartritis periférica en paciente con síndrome de Noonan

RESUMEN

El síndrome de Noonan es un síndrome genético autosómico dominante que presenta una gran variabilidad fenotípica, caracterizado principalmente por dimorfismo facial, cardiopatía congénita y talla baja. Describimos el caso de un paciente de sexo masculino con síndrome de Noonan y espondiloartritis periférica, asociación no descrita en la literatura hasta el momento.

© 2013 Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Noonan's syndrome (NS) is an autosomal dominant genetic syndrome characterized by short stature, cardiac abnormalities, a short neck, chest deformity, a characteristic phenotype with hypertelorism and mental retardation.¹ Its incidence so far has been estimated at 1:1000 to 1:2500 live births.²

The clinical characteristics of patients with NS are a triangular face with a broad forehead, hypertelorism, epicanthus, ptosis, a depressed nasal bridge, micrognathia, small ears, short neck and cardiac abnormalities.^{3,4} Among the musculoskeletal disorders, the most common chest deformities are pectus carinatum and/or pectus excavatum, observed in 70% of cases. In addition there is ulnar

valgus clynobrachydactylia, scoliosis/kyphosis, radioulnar synostosis, hyperextensibility and dental malocclusion.⁵

Other clinical findings are usually bilateral cryptorchidism, which occurs in 60% of male patients, learning and language disorders, attention deficit and depression in 23% of cases.⁶

NS diagnosis is made with clinical findings, according to the criteria formulated by van der Burgt in 1997 and published in 2007, described in Table 1.⁷ Treatment is based on the clinical manifestations.⁸

We report the case of a young male patient with a clinical diagnosis of NS, exhibiting peripheral spondyloarthritis. The relevance of this case is the presentation of a genetic disease with spondyloarthritis, entities that generally do not coexist, allowing us to do a literature review and report this as the first clinical association seen in our region.

Clinical Case

The patient is a 23-year-old male with a family history and diagnosis of NS since the age of 2, manifested by cardiac

[☆] Please cite this article as: Saldarriaga Rivera LM, Fernandes de Melo E, Damião Araujo P, Araujo Silva Filho N, Delgado Quiroz LA, Rios Gomes Bica BE. Presentación de espondiloartritis periférica en paciente con síndrome de Noonan. Reumatol Clin. 2015;11:112–115.

* Corresponding author.

E-mail address: linamarias7@hotmail.com (L.M. Saldarriaga Rivera).

Table 1
Van der Burgt Criteria for the Clinical Diagnosis of Noonan's Syndrome.

Characteristics	Major criteria	Minor criteria	Patient criteria
Facial	Typical face	Suggestive face	Typical face
Cardiac	Pulmonary valve stenosis Hypertrophic cardiomyopathy and/or ECG changes	Other cardiac abnormalities	Intraventricular communication which was treated surgically At the time, no cardiac abnormalities
Height	<3rd percentile	<10th percentile	<3rd percentile
Thoracic	Pectus carinatum and/or pectus excavatum	Width	Pectus carinatum
Family history	First-degree relative diagnosed with Noonan's syndrome	First degree relative with characteristics suggestive of Noonan's syndrome	First degree relative with characteristics suggestive of Noonan's syndrome
Other	Had 3: Mental retardation Cryptorchidism Lymphatic dysplasia	Had one: Mental retardation Cryptorchidism Lymphatic dysplasia	Mental retardation

Diagnosis of Noonan's syndrome:
 -Two major criteria or one major + 2 minor criteria or 3 minor criteria.
 -Typical face with hypertelorism, antimongoloid deviation of palpebral fissures, epicanthus, low-set and rotated ears.

abnormalities (ventricular septal defect diagnosed at 6 months, managed with digoxin, furosemide, antibiotic prophylaxis and surgery until resolved), chest deformity (pectus carinatum), cryptorchidism, bilateral inguinal hernia and surgically corrected phimosis, as well as short stature, a characteristic phenotype with hypertelorism and learning disorders. In 2013 he began a 6 months evolution with ankle, metacarpophalangeal and carpal inflammatory pain, with morning stiffness, and was treated with NSAIDs occasionally with little response; so he was referred to the rheumatology department, with joint pain, bilateral ankle edema and conjunctival hyperemia (Fig. 1).

On physical examination, he had a triangular face with a broad forehead, antimongoloid deviation of palpebral fissures, low-set and rotated ears (Figs. 2 and 3) He presented stunted growth. His height was 1.52 cm (<3rd percentile), weight: 38.4 kg (<3rd percentile), and he presented pectus carinatum, arthritis of the wrist and fourth metacarpophalangeal of the right hand, second and third proximal interphalangeal joint and distally on the left hand, knee as well as the right knee and ankle, with left anserine bursitis and enthesitis of the left Achilles tendon with bilateral plantar fasciitis. The modified Schober test was reduced (2 cm), thoracic expansibility 4 cm, lateral lumbar flexion 9 cm. Occiput-wall distance was 0 cm, tragus-wall 11 cm and finger-floor distance of 37 cm. The patient had no skin, cardiac or genitourinary abnormalities.

Laboratory tests showed: Hb 13.7 mg/dl, Hct 40.8%, WBC: 11,600/mm³; platelets: 456,000/mm³; neutrophils 59%, lymphocytes: 21% CRP: 27.3 mg/L; GOT: 13 U/L; GPT: 24 U/L, GGT: 36 U/L,

**Fig. 1.** Patient with Noonan's syndrome, with bilateral tibial edema.

urea 20 mg/dl, creatinine 0.8 mg/dl, rheumatoid factor: negative, HLA-B27: positive.

X rays of the knees and hands were performed and showed decreased joint space and osteopenia. Magnetic resonance imaging of the lumbosacral spine and hip showed normal vertebral bodies, spinal canal and interapophyseal and sacroiliac joints, with edema of the sacrum. MRI showed edema of the ankles in the posterior region of the calcaneus, thickening and increased density in the insertion of the left Achilles tendon, partial rupture and associated peritendinitis and retrocalcaneal bursitis (Fig. 4). Bone densitometry revealed a lower bone mass than expected for the patients' age.

The ophthalmologic evaluation revealed mild episcleritis, bilateral normal visual acuity, and the patient was treated with ophthalmic corticosteroid 2 times/day.

**Fig. 2.** Patient with Noonan's syndrome phenotype (short stature, short neck, chest deformity, and characteristic triangular face with broad forehead).



Fig. 3. Patient with Noonan's syndrome (short stature, short neck, and chest deformity).



Fig. 4. MRI of the ankle with edema in the posterior region of the calcaneus, thickening and increased density in the insertion of the left Achilles tendon, partial rupture associated to peritendinitis and retrocalcaneal bursitis.

The diagnosis of peripheral spondyloarthritis was made based on the ASAS group classification criteria (arthritis, enthesitis plus HLA-B27 positive).⁹

The BASDAI scale was applied, which measures disease activity in patients with ankylosing spondylitis, with a value of 4. The patient was treated with naproxen 500 mg/12 h sulfasalazine 1.5 g/day, calcium and vitamin D. After 6 months of treatment the patient had a BASDAI of 2, with improvement of clinical symptoms.

Discussion

NS is an uncommon genetic syndrome and an important differential diagnosis in patients with short stature. Musculoskeletal manifestations commonly seen in these patients are pectus

Carinatum type chest deformities, observed in 70% of cases, as well as kyphosis or scoliosis.¹⁰

Bertola et al. described, in a study of 31 patients with NS, musculoskeletal disorders in 27% of the population, the most prevalent being spina bifida and scoliosis, findings not observed in our patient.¹¹

Pozo also described cubitus valgus (50%), clynobrachydactylia (30%), radioulnar synostosis (2%), hyperextensible joints (50%), thoracic and spinal scoliosis in 25% of cases.¹²

None of the studies found arthritis and enthesitis related to the genetic syndrome.

The differential diagnosis of NS includes Turner syndrome, Aarskog syndrome, Klippel-Feil syndrome, fetal alcohol syndrome and primidone embryopathy.¹³

Our patient was diagnosed clinically at 2 years of age and at the time of consultation had major criteria for NS manifested by a typical face, height <3rd percentile, with a first degree family history suggestive of NS, pectus carinatum and absence of cardiac abnormalities associated with the diagnosis of peripheral spondyloarthritis according to the ASAS group criteria.

During initial treatment the patient showed little response to NSAIDs due to their occasional use, taking them only in case of pain. Once the diagnosis of peripheral spondyloarthritis was made and no renal involvement that limited drug use was seen, we started naproxen 500 mg/12 h sulfasalazine 1.5 g/day, calcium and vitamin D with clinical improvement.

After conducting an extensive review of literature in the PubMed and Cochrane databases, we determined that the association of peripheral spondyloarthritis and NS in a young male patient is a relevant fact which no publication to date has documented.

We can also consider that the coexistence of these 2 diseases is a casual relationship and, according to the literature studied, the genetic alterations of NS do not appear to be associated with HLA-B27, which is important for the rheumatologist to know, in addition to the musculoskeletal disorders associated with this genetic syndrome.

Ethical Responsibilities

Protection of people and animals. The authors declare that no experiments have been performed on humans or animals.

Data Privacy. The authors state that no patient data appear in this article.

Right to privacy and informed consent. The authors state that no patient data appear in this article.

Conflict of Interest

The authors declare no conflict of interest.

Financing

None.

References

- Mendez HM, Opitz JM. Noonan syndrome: a review. *Am J Med Genet.* 1985;21:493–506.
- Allanson JE. Noonan syndrome. *J Med Genet.* 1987;24:9–13.
- Ranke MB, Heidemann P, Knupfer C, Enders H, Schmaltz AA, Bierich JR. Noonan syndrome: growth and clinical manifestations in 144 cases. *Eur J Pediatr.* 1988;148:220–7.
- Noonan J, Ehmke D. Associated non cardiac malformations in children with congenital heart disease. *J Pediatr.* 1963;63:468–70.

5. Sharland M, Burch M, Mckenna WM, Patton MA. A clinical study of Noonan syndrome. *Arch Dis Child.* 1992;67:178–83.
6. Ballesta Martínez MJ, Guillén-Navarro E. Síndrome de Noonan. *Protoc Diagn ter Pediatr.* 2010;1:56–63.
7. Van der Burgt I. Noonan syndrome. *Orphanet J Rare Dis.* 2007;2:4.
8. Shaw AC, Kalidas K, Crosby AH, Jeffery S, Patton MA. The natural history of Noonan syndrome: a long-term follow-up study. *Arch Dis Child.* 2007;92:128–32.
9. Rudwaleit M, van der Heijde D, Landewé R, Akkoc N, Brandt J, Chou CT, et al. Criteria the assessment of spondyloarthritis international society classification criteria for peripheral spondyloarthritis and for spondyloarthritis in general. *Ann Rheum Dis.* 2011;70:25–31.
10. Noonan JA. Noonan syndrome and related disorders. *Prog Pediatr Cardiol.* 2005;20:177–85.
11. Bertola DR, Sugayama MS, Albano LM, Gonzalez CH, Chong AK. Noonan syndrome: a clinical and genetic study of 31 patients. *Rev Hosp Clin Fac Med Sao Paulo.* 1999;54:147–50.
12. Pozo JR. Síndrome de Noonan y síndromes relacionados (síndromes neuro-facio-cardio-cutáneos). *Pediatr Integral.* 2010;14:636–47.
13. Opitz JM. The Noonan syndrome. *Am J Med Genet.* 1985;21:515–8.