



Sociedad Española
de Reumatología -
Colegio Mexicano
de Reumatología

Reumatología Clínica

www.reumatologiaclinica.org



Editorial

Glucocorticoids for rheumatoid arthritis in the era of targeted therapies

Glucocorticoides para artritis reumatoide en la era de las terapias dirigidas

Maarten Boers

Epidemiology & Biostatistics, Amsterdam Rheumatology and Immunology Center, Amsterdam University Medical Centers, Vrije Universiteit, Amsterdam, Netherlands

Background/Introduction

Glucocorticoids (GC) constitute one of the cornerstones of therapy for rheumatoid arthritis (RA), and are likely the most frequently prescribed chronic therapy after methotrexate. Both in surveys (QUEST-RA) and in trials about half of RA patients are on chronic GC therapy, with considerable variation between countries.¹ It was expected that the introduction of targeted therapies such as the biologics would lead to reductions in GC use, but this has not really materialized.²

GC are associated with many adverse events, and treatment guidelines always stress that GC use must be as brief as possible, and at the lowest possible dose. So there is a discrepancy between 'the real world' and the guidelines (and pharma marketing).

The purpose of this viewpoint is to provide a realistic perspective on the current state of the art regarding the balance of benefit and harm of GC in RA, and to compare disease-modifying strategies that include GC with strategies that include biologics and synthetic targeted agents. It is not a systematic review, although the reader is pointed to other reviews for reference.

Notes on scope and definitions

Low dose GC therapy is defined as 7.5 mg/d of prednisolone equivalent or less, and intermediate dose as >7.5 and ≤ 30 mg/d.³ Most studies on GC have employed doses between 5 and 10 mg/d. Some have started with a high dose (30–60 mg/d) rapidly 'stepping down' to low dose, and some have looked at 'bridging strategies' where intermediate doses were given for the initial months, either as oral therapy or as repeated intramuscular injections. Intravenous pulse therapy has rarely been studied and falls outside the scope of this viewpoint.

Disease-modifying antirheumatic drugs (DMARDs) are those drugs that are able to alter the course of disease, and this is usually translated to mean that the drug is able to slow or halt the progression of joint damage as assessed on radiographs of hands

and forefeet. Traditional DMARDs such as methotrexate (MTX), sulfasalazine, and hydroxychloroquine were also called 'slow-acting' and distinguished from GC because the latter act rapidly. Until quite recently, GC were not regarded as DMARD because of a wrongly held belief that GC did not slow progression of damage. Despite quite convincing evidence to the contrary (dating back to the early 1960-ies!),⁴ this belief is tenacious, and many review articles on the treatment of RA still distinguish between 'traditional DMARDs' and GC. Newer DMARDs are often called 'Targeted therapies' and include targeted biologics ('bDMARDs') and targeted synthetic agents ('tsDMARDs'; currently only Janus kinase-inhibitors).

'Treat to target' (T2T) is an ill-defined concept: both the specified target and the treatment protocol varies widely; most research in treat-to-target is in early RA.

Regarding outcome measurement, in RA trials ACR (American College of Rheumatology) response (suite: 20-50-70)⁵ and DAS28 (Disease Activity Score-28 joints),⁶ including EULAR response⁷ are most commonly used. Although both perform admirably in distinguishing active from placebo, and also weak from strong drug treatment, they work best in the setting of moderate to severe disease activity. Both are suboptimal in the setting of low disease activity or remission.

Evidence for benefit of treat-to-target and combination therapy in early disease

A systematic review on T2T published in 2017 included 16 studies.⁸ Only 6 of these compared T2T with usual care (usually MTX monotherapy), providing some evidence that T2T is better, via earlier achievement of desired state. The 10 other trials compared different strategies against each other.

There is a substantial body of evidence for the benefit of GC in early RA, but much less for established RA; this is discussed below. This contrasts with the evidence for most other therapies, where the focus is on established RA. Also, most GC trials describe a treatment strategy where GC combined with other DMARDs is compared to DMARDs alone. For non-GC traditional DMARDs there is usually evidence for both combination therapy and

E-mail address: eb@vumc.nl




Fig. 1. Variations on the COBRA schedule. Based on: COBRA: Boers et al.,¹⁹ COBRA-light: den Uyl et al.,²² and COBRA-slim: Verschuereen et al.²³

monotherapy. For the biologics and tsDMARDs, the most frequently studied combination is that with MTX.

In registration (phase 3) trials of new DMARDs, most frequently patients with an insufficient response to MTX are randomized to additional placebo or the new DMARD. This is technically also a combination study, but it supplies little information on the true value of the combination because patients have already shown insufficient response (or failure) on MTX.⁹

For non-GC DMARD strategies mostly initial combo vs MTX monotherapy studies have been performed. Recent and older reviews collected very limited evidence that initial combo is better: most trials were negative, except for some evidence to suggest a benefit of 'triple therapy' (MTX + SSZ + HCQ) over MTX monotherapy.^{10,11}

In the course of their development, all targeted agents have been studied in early RA (MTX naïve patients), where MTX monotherapy was compared with the combination, sometimes also with a third arm (targeted agent monotherapy).

A recent review¹² summarized the results as follows: for disease activity a combination works better than MTX monotherapy; the new agent as monotherapy is only marginally better than MTX monotherapy, with the possible exception of tofacitinib monotherapy. More specifically, MTX + several biologics (abatacept, adalimumab, etanercept, infliximab, rituximab, tocilizumab); and tofacitinib monotherapy resulted in an ACR50 response range of 56–67%, compared to 41% for MTX monotherapy. For damage, several MTX-biologic combinations (adalimumab, etanercept, certolizumab, or infliximab) were statistically superior to MTX monotherapy, but in all treatment groups the estimated mean 1-y change was less than the minimal clinically important difference (5 units on the Sharp van der Heijde score¹³). Since this review, similar evidence for baricitinib has been published.¹⁴

Evidence for GC-DMARD strategies

As stated above, most evidence for the efficacy of GC is in early RA. However, this is about to change as the GLORIA trial has recently closed inclusion and will present results in 2021. It is a large ($n = 452$) pragmatic trial running in 7 European countries that compares the addition of 5 mg/d prednisolone or placebo to standard of care therapy in RA patients aged 65+.¹⁵ Also, a withdrawal study presented at last year's American College of Rheumatology meeting showed that patients in low disease activity on tocilizumab plus low-dose GC were worse off if they tapered the GC, indicating continued efficacy beyond that of tocilizumab.¹⁶

The older literature is already conclusive on the efficacy of GC; effective in this review is shorthand for improvement on signs and symptoms AND slowing of damage progression; effects are consistent, as summarized in Cochrane reviews.^{4,17,18} As designer of COBRA, the first step-down trial, this schedule has a special place in my heart.¹⁹ Since its publication, its evidence has been replicated (BeSt trial^{20,21}) and the schedule has been shown to be effective in lower dose modifications,^{22,23} as shown in Fig. 1. However, more simple schedules are also effective^{24–26}. The benefits of early GC also extend to less starts of expensive biologic therapy,^{20,26} and counteraction of the systemic inflammatory effects that may include glucose intolerance, lipid disorders, mood disturbance and weight loss.^{19,27,28} Finally, GC co-therapy reduces the chance of hypersensitivity reactions to other antirheumatic drugs.¹⁹ In Amsterdam an interesting experiment studied glucose metabolism during a week of 60 or 30 mg/d prednisolone in early untreated RA.²⁹ These patients subsequently went on in the COBRA-light trial.²² 41 patients were given 60 or 30 mg/d. At baseline 23 patients had impaired glucose tolerance, and 3 had unrecognized diabetes type 2! The extent of glucose intolerance correlated with disease activity. After 1 week of either 30 or 60 mg/d prednisolone,




Fig. 2. Systematic review of combination of TNF inhibitors and methotrexate (aTNF + MTX) compared to traditional combination therapy in rheumatoid arthritis. Forest plots show mean (95%CI) difference per trial, and weighted mean difference (random effects model) over all trials. Only traditional combinations (DMARD comb) that contain glucocorticoids (GC, light blue) are as good as aTNF + MTX, other combinations (dark blue) are not. Middle blue rhomboid: overall estimate. DAS28: Disease activity Score-28 joints; SvdH (erosion): Sharp van der Heijde radiographic damage erosion subscore; Double: MTX + sulfasalazine; Triple: Double + hydroxychloroquine. GC: glucocorticoids; iv: intravenous. Trial references: APPEAL,³⁹ BeSt (T is trial arm 3, with COBRA; D is trial arm 2, with Double)²¹; IMPROVED⁴⁰; LARA⁴¹; RACAT⁴²; TEAR⁴³; IDEA⁴⁴; SWEFOT.⁴⁵ Adapted from Graudal et al.¹⁰ with permission.

the mean glucose tolerance was unchanged. However, 7 patients with initially impaired tolerance progressed to diabetes mellitus, whereas 9 regressed to normal! Even in these early patients, progression to diabetes was more likely with longer symptom duration (36 v 16 weeks).

So how does combination therapy that includes GC compare with biologic therapy? Quite well, actually. The most well-known example is the BeSt trial where infliximab plus high dose methotrexate was indistinguishable from traditional COBRA therapy (treatment schedule, see Fig. 1) in efficacy.²⁰ A recent review confirmed this with other GC schedules and biologics (Fig. 2).¹⁰ These results strongly suggest biologics should not be started in early RA before a combination that includes GC has been tried.

Harm

Most concern with GC in RA is not whether they work, but the 'certainty' that the harm outweighs any benefit. The number of observational studies that 'prove' this point increases every day, but in this (and many other) setting observational studies can never overcome the strong confounding by indication.^{30–32} This occurs because patients with poor prognosis are not only more likely to receive these agents, and poor prognosis RA is itself associated with many of the adverse events that are traditionally associated with GC. A recent example of this, only presented in abstract is a study

from the CATCH cohort, that looked at 1689 early RA patients³³: 59% received GC within the first 3 months (30% oral). These patients were older, less likely to be employed, had shorter disease duration, were more often rheumatoid factor or anti-citrullinated protein antibody positive, and were worse in all indicators of disease activity, and in physical disability scores. They had a worse course in the first year including more biologic starts.

In contrast, the experience from randomized trials (where by definition GC therapy is not associated with RA prognosis) consistently contradicts that low-dose GC (≤ 7.5 mg prednisolone equivalent/d) given for extended periods is associated with excessive harm. However such evidence has proved hard to publish: a systematic review is still languishing in abstract form.³⁴ However, a published case in point is our series on the long-term effects seen in the COBRA cohort: evidence for prolonged protection against damage,³⁵ and no signs of excessive harm³⁶ or increased mortality.³⁷ It is certain that GC can cause serious harm, but it is highly likely that such harm is mostly seen in conditions where GC are given for long periods in high doses. Such conditions include complicated lupus and vasculitis, but most are non-rheumatologic, such as lymphoma.

In conclusion, it is time we dropped our strange reservations against the inclusion of GC in our treatment armamentarium, as a 'normal' drug, with pro's and con's like any other treatment. This includes getting rid of our collective self-denial, where guidelines ('short-term', 'taper as rapidly as possible'³⁸) are completely at odds with daily practice (about 50% of our RA patients on long-term GC therapy), but also optimal attention to effective preventive measures (screening and treating comorbidity, bone protection). Finally, more public funding and research effort must be spent on acquiring more knowledge about this essential agent.

References

- Sokka T, Kautiainen H, Toloza S, Makinen H, Verstappen SM, Lund Hetland M, et al. QUEST-RA: quantitative clinical assessment of patients with rheumatoid arthritis seen in standard rheumatology care in 15 countries. *Ann Rheum Dis.* 2007;66:1491–6.
- Makol A, Davis JM, 3rd, Crowson CS, Thorneau TM, Gabriel SE, Matteson EL. Time trends in glucocorticoid use in rheumatoid arthritis: results from a population-based inception cohort, 1980–1994 versus 1995–2007. *Arthritis Care Res (Hoboken).* 2014;66:1482–8.
- Buttgereit F, Da Silva JA, Boers M, Burmester GR, Cutolo M, Jacobs J, et al. Standardised nomenclature for glucocorticoid dosages and glucocorticoid treatment regimens: current questions and tentative answers in rheumatology. *Ann Rheum Dis.* 2002;61:718–22.
- Kirwan JR, Bijlsma JW, Boers M, Shea BJ. Effects of glucocorticoids on radiological progression in rheumatoid arthritis. *Cochrane Database Syst Rev.* 2007. CD006356.
- Felson DT, Anderson JJ, Boers M, Bombardier C, Furst D, Goldsmith C, et al. American College of Rheumatology Preliminary definition of improvement in rheumatoid arthritis. *Arthritis Rheum.* 1995;38:727–35.
- Prevoo ML, van't Hof MA, Kuper HH, van Leeuwen MA, van de Putte LB, van Riel PL. Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum.* 1995;38:44–8.
- van Gestel AM, Prevoo ML, van't Hof MA, van Rijswijk MH, van de Putte LB, van Riel PL. Development and validation of the European League Against Rheumatism response criteria for rheumatoid arthritis. Comparison with the preliminary American College of Rheumatology and the World Health Organization/International League Against Rheumatism Criteria. *Arthritis Rheum.* 1996;39:34–40.
- Wailoo A, Hock ES, Stevenson M, Martyn-St James M, Rawdin A, Simpson E, et al. The clinical effectiveness and cost-effectiveness of treat-to-target strategies in rheumatoid arthritis: a systematic review and cost-effectiveness analysis. *Health Technol Assess.* 2017;21:1–258.
- Boers M. Add-on or step-up trials for new drug development in rheumatoid arthritis: a new standard? *Arthritis Rheum.* 2003;48:1481–3.
- Graudal N, Hubeck-Graudal T, Faurschou M, Baslund B, Jürgens G. Combination therapy with and without tumor necrosis factor inhibitors in rheumatoid arthritis: a meta-analysis of randomized trials. *Arthritis Care Res.* 2015;67:1487–95.
- Verhoeven AC, Boers M, Tugwell P. Combination therapy in rheumatoid arthritis – updated systematic review. *Br J Rheumatol.* 1998;37:612–9.
- Hazlewood GS, Barnabe C, Tomlinson G, Marshall D, Devove DJ, Bombardier C. Methotrexate monotherapy and methotrexate combination therapy with

- traditional and biologic disease modifying anti-rheumatic drugs for rheumatoid arthritis: a network meta-analysis. *Cochrane Database Syst Rev.* 2016. CD010227.
13. Bruynesteyn K, van Der Heijde D, Boers M, Saudan A, Peloso P, Paulus H, et al. Determination of the minimal clinically important difference in rheumatoid arthritis joint damage of the Sharp/van der Heijde and Larsen/Scott scoring methods by clinical experts and comparison with the smallest detectable difference. *Arthritis Rheum.* 2002;46:913–20.
 14. van der Heijde D, Durez P, Schett G, Naredo E, Ostergaard M, Meszaros G, et al. Structural damage progression in patients with early rheumatoid arthritis treated with methotrexate, baricitinib, or baricitinib plus methotrexate based on clinical response in the phase 3 RA-BEGIN study. *Clin Rheumatol.* 2018;37:2381–90.
 15. Hartman L, Rasch LA, Klausch T, Bijlsma HWJ, Christensen R, Smulders YM, et al. Harm, benefit and costs associated with low-dose glucocorticoids added to the treatment strategies for rheumatoid arthritis in elderly patients (GLORIA trial): study protocol for a randomised controlled trial. *Trials.* 2018;19:67.
 16. Burmester GR, Buttgereit F, Bernasconi C, Alvaro-Gracia JM, Castro N, Dougados M, et al. A randomized controlled 24-week trial evaluating the safety and efficacy. *Arthritis Rheumatol.* 2018;70 Suppl. 10.
 17. Gotsche PC, Johansen HK. Short-term low-dose corticosteroids vs placebo and nonsteroidal antiinflammatory drugs in rheumatoid arthritis. *Cochrane Database Syst Rev.* 2000. CD000189.
 18. Criswell LA, Saag KG, Sems KM, Welch V, Shea B, Wells G, et al. Moderate-term, low-dose corticosteroids for rheumatoid arthritis. *Cochrane Database Syst Rev.* 2000. CD001158.
 19. Boers M, Verhoeven AC, Markuse HM, van de Laar MA, Westhovens R, van Denderen JC, et al. Randomised comparison of combined step-down prednisolone, methotrexate and sulphasalazine with sulphasalazine alone in early rheumatoid arthritis [see comments] [published erratum appears in *Lancet* 1998 Jan 17;351(9097):220]. *Lancet.* 1997;350:309–18.
 20. Goekoop-Ruiterman YP, de Vries-Bouwstra JK, Allaart CF, van Zeben D, Kerstens PJ, Hazes JM, et al. Comparison of treatment strategies in early rheumatoid arthritis: a randomized trial. *Ann Intern Med.* 2007;146:406–15.
 21. Goekoop-Ruiterman YP, de Vries-Bouwstra JK, Allaart CF, van Zeben D, Kerstens PJ, Hazes JM, et al. Clinical and radiographic outcomes of four different treatment strategies in patients with early rheumatoid arthritis (the BeSt study): a randomized, controlled trial. *Arthritis Rheum.* 2005;52:3381–90.
 22. den Uyl D, ter Wee M, Boers M, Kerstens P, Voskuyl A, Nurmohamed M, et al. A non-inferiority trial of an attenuated combination strategy ('COBRA-light') compared to the original COBRA strategy: clinical results after 26 weeks. *Ann Rheum Dis.* 2014;73:1071–8.
 23. Verschueren P, De Cock D, Corluy L, Joos R, Langenaken C, Taelman V, et al. Effectiveness of methotrexate with step-down glucocorticoid remission induction (COBRA Slim) versus other intensive treatment strategies for early rheumatoid arthritis in a treat-to-target approach: 1-year results of CareRA, a randomised pragmatic open-label superiority trial. *Ann Rheum Dis.* 2017;76:511–20.
 24. Kirwan JR, for the arthritis rheumatism council low-dose glucocorticosteroid study group. The effects of glucocorticosteroids on joint destruction in rheumatoid arthritis. *N Engl J Med.* 1995;333:142–6.
 25. Möttönen T, Hannonen P, Leirisalo-Repo M, Nissila M, Kautiainen H, Korpela M, et al. Comparison of combination therapy with single-drug therapy in early rheumatoid arthritis: a randomised trial. *Lancet.* 1999;353:1568–73.
 26. Bakker MF, Jacobs JW, Welsing PM, Verstappen SM, Tekstra J, Ton E, et al. Low-dose prednisone inclusion in a methotrexate-based, tight control strategy for early rheumatoid arthritis: a randomized trial. *Ann Internal Med.* 2012;156:329–39.
 27. Hoes JN, van der Goes MC, van Raalte DH, van der Zijl NJ, den Uyl D, Lems WF, et al. Glucose tolerance, insulin sensitivity and beta-cell function in patients with rheumatoid arthritis treated with or without low-to-medium dose glucocorticoids. *Ann Rheum Dis.* 2011;70:1887–94.
 28. Boers M, Nurmohamed MT, Doelman CJ, Lard LR, Verhoeven AC, Voskuyl AE, et al. Influence of glucocorticoids and disease activity on total and high density lipoprotein cholesterol in patients with rheumatoid arthritis. *Ann Rheum Dis.* 2003;62:842–5.
 29. den Uyl D, van Raalte DH, Nurmohamed MT, Lems WF, Bijlsma JW, Hoes JN, et al. Metabolic effects of high-dose prednisolone treatment in early rheumatoid arthritis: balance between diabetogenic effects and inflammation reduction. *Arthritis Rheum.* 2012;64:639–46.
 30. Boers M. Studying the benefit/risk ratio of glucocorticoids in rheumatoid arthritis. *J Rheumatol.* 2007;34:661–3.
 31. Da Silva JA, Jacobs JW, Kirwan JR, Boers M, Saag KG, Ines LB, et al. Safety of low dose glucocorticoid treatment in rheumatoid arthritis: published evidence and prospective trial data. *Ann Rheum Dis.* 2006;65:285–93.
 32. Boers M. Bias in its coat of many colors. *J Rheumatol.* 2001;28:1747–8.
 33. Andersen K, Lin D, Bartlett SJ, Boire G, Haraoui B, Hitchon C, et al. Characteristics and outcomes associated with early corticosteroid use in a large multicenter Canadian RA Cohort [abstract]. *Arthritis Rheumatol.* 2015;67 Suppl. 10.
 34. Tarp S, Bartels EM, Kirwan JR, et al. Adverse effects of glucocorticoid therapy in rheumatoid arthritis: a systematic review and meta-analysis of randomized controlled trials. *Ann Rheum Dis.* 2012;71 Suppl. 3:207.
 35. Landewe RB, Boers M, Verhoeven AC, Westhovens R, van De Laar MA, Markuse HM, et al. COBRA combination therapy in patients with early rheumatoid arthritis: Long-term structural benefits of a brief intervention. *Arthritis Rheum.* 2002;46:347–56.
 36. van Tuyl LH, Boers M, Lems WF, Landewe RB, Han H, van der Linden S, et al. Survival, comorbidities and joint damage 11 years after the COBRA combination therapy trial in early rheumatoid arthritis. *Ann Rheum Dis.* 2010;69:807–12.
 37. Poppelaars PB, van Tuyl LHD, Boers M. Normal mortality of the COBRA early rheumatoid arthritis trial cohort after 23 years of follow-up. *Ann Rheum Dis.* 2019;78:586–9.
 38. Smolen JS, Landewe R, Bijlsma J, Burmester G, Chatzidionysiou K, Dougados M, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis.* 2017;76:960–77.
 39. Kim HY, Hsu PN, Barba M, Sulaiman W, Robertson D, Vlahos B, et al. Randomized comparison of etanercept with usual therapy in an Asian population with active rheumatoid arthritis: the APPEAL trial. *Int J Rheum Dis.* 2012;15:188–96.
 40. Heimans L, Wevers-de Boer KV, Visser K, Goekoop RJ, van Oosterhout M, Harbers JB, et al. A two-step treatment strategy trial in patients with early arthritis aimed at achieving remission: the IMPROVED study. *Ann Rheum Dis.* 2014;73:1356–61.
 41. Machado DA, Guzman RM, Xavier RM, Simon JA, Mele L, Pedersen R, et al. Open-label observation of addition of etanercept versus a conventional disease-modifying antirheumatic drug in subjects with active rheumatoid arthritis despite methotrexate therapy in the Latin American region. *J Clin Rheumatol.* 2014;20:25–33.
 42. O'Dell JR, Mikuls TR, Taylor TH, Ahluwalia V, Brophy M, Warren SR, et al. Therapies for active rheumatoid arthritis after methotrexate failure. *N Engl J Med.* 2013;369:307–18.
 43. O'Dell JR, Leff R, Paulsen G, Haire C, Mallek J, Eckhoff PJ, et al. Treatment of rheumatoid arthritis with methotrexate and hydroxychloroquine, methotrexate and sulfasalazine, or a combination of the three medications: results of a two-year, randomized, double-blind, placebo-controlled trial. *Arthritis Rheum.* 2002;46:1164–70.
 44. Nam JL, Villeneuve E, Hensor EM, Conaghan PG, Keen HI, Buch MH, et al. Remission induction comparing infliximab and high-dose intravenous steroid, followed by treat-to-target: a double-blind, randomised, controlled trial in new-onset, treatment-naïve, rheumatoid arthritis (the IDEA study). *Ann Rheum Dis.* 2014;73:75–85.
 45. van Vollenhoven RF, Geborek P, Forslind K, Albertsson K, Ernestam S, Petersson IF, et al. Conventional combination treatment versus biological treatment in methotrexate-refractory early rheumatoid arthritis: 2 year follow-up of the randomised, non-blinded, parallel-group Swefot trial. *Lancet.* 2012;379:1712–20.