



Sociedad Española
de Reumatología -
Colegio Mexicano
de Reumatología

Reumatología Clínica

www.reumatologiaclinica.org



Images in Clinical Rheumatology

Arthralgias and Articular Hyperlaxitude in Women with Ophthalmopathy and Early Deafness[☆]



Artralgias e hiperlaxitud articular en mujer con oftalmopatía y sordera precoz

Francisco Gallo Puelles,^{a,*} Ana Teresa Serrano Anton,^{b,c} Vanesa Lopez Gonzalez,^{b,c} Jose Hurtado Martinez^d

^a Servicio de Reumatología, Hospital Clínico Universitario Virgen de la Arrixaca, El Palmar, Murcia, Spain

^b Sección de Genética Médica, Servicio de Pediatría, Hospital Clínico Universitario Virgen de la Arrixaca, IMIB-Arrixaca, El Palmar, Murcia, Spain

^c CIBERER-ISCIII, Spain

^d Servicio de Cardiología, Hospital Clínico Universitario Virgen de la Arrixaca, El Palmar, Murcia, Spain

ARTICLE INFO

Article history:

Received 16 October 2018

Accepted 16 October 2018

Available online 9 December 2019

Clinical case

A twenty-six-year-old woman, with arthralgia and crepitation in her hands, feet and knees, and longstanding camptodactyly.^{1,2}

Personal history: bilateral myopia, retinal vitreous degeneration and sensorineural hearing loss since childhood. **Family history:** polyarthralgia, ophthalmological anomalies (retinal vitreous degeneration, retinal detachment, myopia and cataracts), hearing loss, scoliosis and similar phenotype in the paternal line^{3,4} (Fig. 1). Physical examination revealed marfanoid habitus, standard size, scoliosis, joint hyperlaxity (non-cutaneous),⁵ particular facial features (mediofacial hypoplasia, ocular proptosis, depressed nasal bridge, macrognathia)⁶ (Fig. 2A and B) and elongated fingers and toes with camptodactyly⁷ (Fig. 2D).

Diagnosis/progress

Blood tests: ESR 10 mm/1st hour, CRP .3 (<.5 mg/dl); normal haemogram and TSH/T4L, RF/negative autoimmunity (ANA, ENA, ANCA, ACPA). **Radiology:** scoliosis, increased metacarpal length (Fig. 2C). Normal knee MRI and echocardiography.

Given the possibility of hereditary collagenopathy we contacted the medical genetics department, who suspected Stickler syndrome, autosomal disease (AD), dominant and progressive.^{1,9} A molecular study of implicated genes was requested: COL2A1, COL11A1 and COL11A2,^{3,4,8–10} by means of next-generation sequencing (NGS) panel with detection of pathogenic heterozygous variant (not described) c.598C>T (p.Gln200) of the COL2A1 gene. Compatible familial segregation analysis (parents/siblings).^{3,8–10}

[☆] Please cite this article as: Gallo Puelles F, Serrano Anton AT, Lopez Gonzalez V, Hurtado Martinez J. Artralgias e hiperlaxitud articular en mujer con oftalmopatía y sordera precoz. Reumatol Clin. 2020;16:423–25.

* Corresponding author.

E-mail address: fgallopuelles@hotmail.com (F. Gallo Puelles).




Fig. 1. Family tree or genogram. Paternal family line affected. Autosomal dominant inheritance.

Discussion

Stickler syndrome (OMIM: 108300) is a hereditary and progressive connective tissue disorder characterised by skeletal abnormalities (arthralgia, joint hyperlaxity and scoliosis), ocular abnormalities (early-onset myopia, strabismus, cataract and retinal detachment), craniofacial abnormalities (midface hypoplasia, broad/flattened nasal bridge, cleft palate and macro/retrognathia) and sensorineural hearing loss.^{2,5–8,10} With an estimated

incidence of 1/7500, this is a rare congenital collagenopathy, caused by mutation of the COL2A1 AD gene, of variable expressivity, even within the family.^{7–10} In spite of the time available in consultations, it is essential to take an adequate history in our routine practice. A personal and family history of ophthalmopathy and deafness, together with close collaboration with medical genetics were determining factors in the correct diagnosis, prognosis and final treatment of the patient and her family.



Fig. 2. Frontal phenotype (A), lateral (B), hand radiology (C) and arachnodactyly (D). Depressed nasal bridge, face hypoplasia, macrognathia, proptosis, low implantation of the auricular pavilion. Elongated hands and hyperlaxity. Increased length of metacarpals and phalanges.

Conflict of interests

The authors have no conflict of interests to declare.

Acknowledgement

The authors would like to express their gratitude for being able to collaborate in your journal and disseminate their article.

References

- De Wergifosse I, Westhovens R. Osteoarthritis at young age, a diagnostic challenge: a case of Stickler syndrome. *Open Rheumatol J.* 2014;8:100–2.
- Li K, Thorne C. Adult presentation of Stickler syndrome type III. *Clin Rheumatol.* 2010;29:795–7.
- Guo L, Elcioglu NH, Wang Z, Demirkol YK, Isguven P, Matsumoto N, et al. Novel and recurrent *COL11A1* and *COL2A1* mutations in the Marshall–Stickler syndrome spectrum. *Hum Genome Var.* 2017;4:17040.
- Avcin T, Makitie O, Susic M, Miller S, Thorne C, Tenenbaum J, et al. Early-onset osteoarthritis due to otospondylomegapiphyseal dysplasia in a family with a novel splicing mutation of the *COL11A2* gene. *J Rheumatol.* 2008;35:920–6.
- Cattalini M, Khubchandani R, Cimaz R. When flexibility is not necessarily a virtue: a review of hypermobility syndromes and chronic or recurrent musculoskeletal pain in children. *Pediatr Rheumatol Online J.* 2015;13:40.
- Acke FR, Dhooge IJ, Malfait F, de Leenheer EM, de Pauw GA, et al. Cephalometrics in Stickler syndrome: objectification of the typical facial appearance. *J Craniomaxillofac Surg.* 2016;44:848–53.